

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

10667187  
12.17.03



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 401/00</b>		A2	(11) International Publication Number: <b>WO 98/52937</b> (43) International Publication Date: 26 November 1998 (26.11.98)
(21) International Application Number: PCT/US98/10807 (22) International Filing Date: 22 May 1998 (22.05.98)  (30) Priority Data: 60/047,535 22 May 1997 (22.05.97) US  (71) Applicant (for all designated States except US): G.D. SEARLE AND CO. [US/US]; P.O. Box 5110, Chicago, IL 60680 (US).  (72) Inventors; and (75) Inventors/Applicants (for US only): ANANTANARAYAN, Ashok [US/US]; 54 Lisk Drive, Hainesville, IL 60015 (US). CLARE, Michael [GB/US]; 5154 West Brown Street, Skokie, IL 60077 (US). GENG, Lifeng [CN/US]; 5300 Davis Street, Skokie, IL 60077 (US). HANSON, Gunnar, J. [US/US]; 7410 Keystone Avenue, Skokie, IL 60076 (US). PARTIS, Richard, A. [US/US]; 2221 Noyes Street, Evanston, IL 60012 (US). STEALEY, Michael, A. [US/US]; 502 Juniper Parkway, Libertyville, IL 60048 (US). WEIER, Richard, M. [US/US]; 240 Hickory Court, Lake Bluff, IL 60044 (US).			(74) Agents: WILLIAMS, Scott, A. et al.; Senniger, Powers, Leavitt & Roedel, 16th floor Fl, One Metropolitan Square, St. Louis, MO 63102 (US).  (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published Without international search report and to be republished upon receipt of that report.
(54) Title: 3(5)-HETEROARYL SUBSTITUTED PYRAZOLES AS p38 KINASE INHIBITORS			
(57) Abstract  A class of pyrazole derivatives is described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula (I), wherein R <sup>1</sup> , R <sup>2</sup> , Ar <sup>1</sup> and HetAr <sup>2</sup> are as described in the specification.			
<div style="text-align: center;"><p>(I)</p></div>			

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

3(5)-HETEROARYL SUBSTITUTED PYRAZOLES  
AS p38 KINASE INHIBITORS

5 Cross-Reference to Related Application

This application claims priority from U.S.  
Provisional Application Serial No. 60/047,535 filed May  
22, 1997.

10

Field of the Invention

This invention relates to a novel group of pyrazole  
compounds, compositions and methods for treating p38  
15 kinase mediated disorders.

Background of the Invention

Mitogen-activated protein kinases (MAP) is a family  
20 of proline-directed serine/threonine kinases that  
activate their substrates by dual phosphorylation. The  
kinases are activated by a variety of signals including  
nutritional and osmotic stress, UV light, growth factors,  
endotoxin and inflammatory cytokines. The p38 MAP kinase  
25 group is a MAP family of various isoforms, including  
p38 $\alpha$ , p38 $\beta$  and p38 $\gamma$ , and is responsible for  
phosphorylating and activating transcription factors  
(e.g. ATF2, CHOP and MEF2C) as well as other kinases  
(e.g. MAPKAP-2 and MAPKAP-3). The p38 isoforms are  
30 activated by bacterial lipopolysaccharide, physical and  
chemical stress and by pro-inflammatory cytokines,  
including tumor necrosis factor (TNF- $\alpha$ ) and interleukin-1  
(IL-1). The products of the p38 phosphorylation mediate  
the production of inflammatory cytokines, including TNF  
35 and IL-1, and cyclooxygenase-2.

TNF- $\alpha$  is a cytokine produced primarily by activated  
monocytes and macrophages. Excessive or unregulated TNF

production has been implicated in mediating a number of diseases. Recent studies indicate that TNF has a causative role in the pathogenesis of rheumatoid arthritis. Additional studies demonstrate that  
5 inhibition of TNF has broad application in the treatment of inflammation, inflammatory bowel disease, multiple sclerosis and asthma.

TNF has also been implicated in viral infections, such as HIV, influenza virus, and herpes virus including  
10 herpes simplex virus type-1 (HSV-1), herpes simplex virus type-2 (HSV-2), cytomegalovirus (CMV), varicella-zoster virus (VZV), Epstein-Barr virus, human herpesvirus-6 (HHV-6), human herpesvirus-7 (HHV-7), human herpesvirus-8 (HHV-8), pseudorabies and rhinotracheitis, among others.

15 IL-8 is another pro-inflammatory cytokine, which is produced by mononuclear cells, fibroblasts, endothelial cells, and keratinocytes, and is associated with conditions including inflammation.

IL-1 is produced by activated monocytes and  
20 macrophages and is involved in the inflammatory response. IL-1 plays a role in many pathophysiological responses including rheumatoid arthritis, fever and reduction of bone resorption.

TNF, IL-1 and IL-8 affect a wide variety of cells  
25 and tissues and are important inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these cytokines by inhibition of the p38 kinase is of benefit in controlling, reducing and alleviating many of these disease states.

30 Various pyrazoles have previously been described. U.S. Patent No. 4,000,281, to Beiler and Binon, describes 4,5-aryl, heteroaryl substituted pyrazoles with antiviral activity against both RNA and DNA viruses such as myxoviruses, adenoviruses, rhinoviruses, and various  
35 viruses of the herpes group. WO 92/19615, published November 12, 1992, describes pyrazoles as novel

fungicides. U. S. Patent No. 3,984,431, to Cueremy and Renault, describes derivatives of pyrazole-5-acetic acid as having antiinflammatory activity. Specifically, [1-isobutyl-3,4-diphenyl-1H-pyrazol-5-yl]acetic acid is described. U. S. Patent No. 3,245,093 to Hinsgen et al, describes a process for preparing pyrazoles. WO 83/00330, published February 3, 1983, describes new process for the preparation of diphenyl-3,4-methyl-5-pyrazole derivatives. WO 95/06036, published for preparing pyrazole and its derivatives. U.S. patent 5,589,439, to T. Goto, et al., describes tetrazole derivatives and their use as herbicides. EP 515041 describes pyrimidyl substituted pyrazole derivatives as novel agricultural fungicides. Japanese Patent 4,145,081 describes pyrazolecarboxylic acid derivatives as herbicides used in paddy fields, dry fields as well as non-agricultural areas. Japanese Patent 5,345,772 describes novel pyrazole derivatives having potent inhibitory activity against acetylcholinesterase.

Pyrazoles have been described for use in the treatment of inflammation. Japanese Patent 5,017,470 describes synthesis of pyrazole derivatives as anti-inflammatory, anti-rheumatic, anti-bacterial and anti-viral drugs. EP 115640, published December 30, 1983, describes 4-imidazolyl-pyrazole derivatives as inhibitors of thromboxane synthesis. 3-(4-Isopropyl-1-methylcyclohex-1-yl)-4-(imidazol-1-yl)-1H-pyrazole is specifically described. WO 97/01551, published January 16, 1997, describes pyrazole compounds as adenosine antagonists. 4-(3-Oxo-2,3-dihydropyridazin-6-yl)-3-phenylpyrazole is specifically described. U.S. Patent No. 5,134,142, to Matsuo et al. describes 1,5-diaryl pyrazoles as having anti-inflammatory activity.


U.S. Patent No. 5,559,137 to Adams et al, describes novel pyrazoles (1,3,4,-substituted) as inhibitors of cytokines used in the treatment of cytokine diseases.

Specifically, 3-(4-fluorophenyl)-1-(4-methylsulfinylphenyl)-4-(4-pyridyl)-5H-pyrazole is described. WO 96/03385, published February 8, 1996, describes 3,4-substituted pyrazoles, as having anti-inflammatory activity. Specifically, 4-[1-ethyl-4-(4-pyridyl)-5-trifluoromethyl-1H-pyrazol-3-yl]benzenesulfonamide is described.

The invention's pyrazolyl compounds are found to show usefulness as p38 kinase inhibitors.

### Description of the Invention

A class of substituted pyrazolyl compounds useful in treating p38 mediated disorders is defined by Formula I:



wherein

R<sup>1</sup> is selected from hydrido, alkyl, cycloalkyl, alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene, haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl, mercaptoalkyl, alkylthioalkylene, amino, alkylamino, arylamino, aminoalkyl, alkylaminoalkylene, heterocyclylalkylene, aminocarbonylalkylene, and alkylaminocarbonylalkylene; and

R<sup>2</sup> is selected from hydrido, alkyl, alkenyl, alkynyl, heterocyclyl, haloalkyl, heterocyclylalkyl, amino, alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy, alkoxycarbonyl, carboxyalkyl, aminocarbonylamino,

alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl, alkylsulfonylamino, aminosulfonylamino, alkylaminosulfonylamino, and alkynylamino; wherein the heterocyclyl and heterocyclylalkyl groups are optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl, haloalkyl, amino, cyano, and hydroxy; and

Ar<sup>1</sup> is aryl optionally substituted with one or more radicals independently selected from halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino, aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl; and

HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

Compounds of Formula I would be useful for, but not limited to, the treatment of any disorder or disease state in a human, or other mammal, which is exacerbated or caused by excessive or unregulated TNF or p38 kinase production by such mammal. Accordingly, the present invention provides a method of treating a cytokine-mediated disease which comprises administering an effective cytokine-interfering amount of a compound of Formula I, or a pharmaceutically acceptable salt or tautomer thereof.



Compounds of Formula I would be useful for, but not limited to, the treatment of inflammation in a subject, and for use as antipyretics for the treatment of fever. Compounds of the invention would be useful to treat

5 arthritis, including but not limited to, rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis, osteoarthritis, gouty arthritis and other arthritic conditions. Such compounds would be useful for

10 the treatment of pulmonary disorders or lung inflammation, including adult respiratory distress syndrome, pulmonary sarcoidosis, asthma, silicosis, and chronic pulmonary inflammatory disease. The compounds are also useful for the treatment of viral and bacterial

15 infections, including sepsis, septic shock, gram negative sepsis, malaria, meningitis, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), pneumonia, and herpesvirus. The

20 compounds are also useful for the treatment of bone resorption diseases, such as osteoporosis, endotoxic shock, toxic shock syndrome, reperfusion injury, autoimmune disease including graft vs. host reaction and allograft rejections, cardiovascular diseases including

25 atherosclerosis, thrombosis, congestive heart failure, and cardiac reperfusion injury, renal reperfusion injury, liver disease and nephritis, and myalgias due to infection.

The compounds are also useful for the treatment of

30 influenza, multiple sclerosis, cancer, diabetes, systemic lupus erythematosus (SLE), skin-related conditions such as psoriasis, eczema, burns, dermatitis, keloid formation, and scar tissue formation. Compounds of the invention also would be useful to treat gastrointestinal

35 conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome and



ulcerative colitis. The compounds would also be useful in the treatment of ophthalmic diseases, such as retinitis, retinopathies, uveitis, ocular photophobia, and of acute injury to the eye tissue. Compounds of the invention also would be useful for treatment of angiogenesis, including neoplasia; metastasis; ophthalmological conditions such as corneal graft rejection, ocular neovascularization, retinal neovascularization including neovascularization following injury or infection, diabetic retinopathy, retrolental fibroplasia and neovascular glaucoma; ulcerative diseases such as gastric ulcer; pathological, but non-malignant, conditions such as hemangiomas, including infantile hemangiomas, angiofibroma of the nasopharynx and avascular necrosis of bone; diabetic nephropathy and cardiomyopathy; and disorders of the female reproductive system such as endometriosis. The compounds of the invention may also be useful for preventing the production of cyclooxygenase-2.

Besides being useful for human treatment, these compounds are also useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.

The present compounds may also be used in co-therapies, partially or completely, in place of other conventional antiinflammatories, such as together with steroids, cyclooxygenase-2 inhibitors, NSAIDs, DMARDs, immunosuppressive agents, 5-lipoxygenase inhibitors, LTB<sub>4</sub> antagonists and LTA<sub>4</sub> hydrolase inhibitors.

As used herein, the term "TNF mediated disorder" refers to any and all disorders and disease states in which TNF plays a role, either by control of TNF itself, or by TNF causing another monokine to be released, such as but not limited to IL-1, IL-6 or IL-8. A disease state in which, for instance, IL-1 is a major component, and

whose production or action, is exacerbated or secreted in response to TNF, would therefore be considered a disorder mediated by TNF.

As used herein, the term "p38 mediated disorder" refers to any and all disorders and disease states in which p38 plays a role, either by control of p38 itself, or by p38 causing another factor to be released, such as but not limited to IL-1, IL-6 or IL-8. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to p38, would therefore be considered a disorder mediated by p38.

As TNF- $\beta$  has close structural homology with TNF- $\alpha$  (also known as cachectin), and since each induces similar biologic responses and binds to the same cellular receptor, the synthesis of both TNF- $\alpha$  and TNF- $\beta$  are inhibited by the compounds of the present invention and thus are herein referred to collectively as "TNF" unless specifically delineated otherwise.

A preferred class of compounds consists of those compounds of Formula I wherein

R<sup>1</sup> is selected from hydrido, lower alkyl, lower cycloalkyl, lower cycloalkylalkylene, lower haloalkyl, lower hydroxyalkyl, lower alkenyl, lower alkynyl, lower heterocyclyl, lower aralkyl, lower alkoxyalkyl, lower mercaptoalkyl, lower alkylthioalkylene, amino, lower alkylamino, lower arylamino, lower aminoalkyl, lower alkylaminoalkylene, lower heterocyclylalkylene, lower aminocarbonylalkylene, and lower alkylaminocarbonylalkylene; and

R<sup>2</sup> is selected from hydrido, lower alkyl, lower alkenyl, lower alkynyl, lower haloalkyl, lower heterocyclyl, lower heterocyclylalkylene, amino, lower alkylamino, lower alkynylamino, lower aminoalkyl, lower alkylthio, lower carboxy, lower alkoxycarbonyl, lower carboxyalkyl, lower aminocarbonylamino, lower

alkylaminocarbonylamino, lower alkylsulfonyl, lower  
aminosulfonyl, lower alkylsulfonylamino, lower  
aminosulfonylamino, and lower alkylaminosulfonylamino,  
wherein the heterocyclyl and heterocyclylalkyl groups are  
5 optionally substituted with one or more radicals  
independently selected from lower alkylthio, lower  
alkylsulfonyl, lower alkylsulfinyl, halo, lower alkyl,  
lower alkoxy, aryloxy, lower heterocyclyl, lower  
haloalkyl, amino, and cyano; and

10 Ar<sup>1</sup> is selected from phenyl, biphenyl, and naphthyl,  
wherein Ar<sup>1</sup> is optionally substituted with one or more  
radicals independently selected from lower alkylthio,  
lower alkylsulfonyl, aminosulfonyl, halo, lower alkyl,  
lower alkenyl, lower alkynyl, lower alkylsulfinyl, cyano,  
15 lower alkoxycarbonyl, aminocarbonyl, formyl, lower  
alkylcarbonylamino, lower haloalkyl, lower alkoxy, lower  
alkenyloxy, lower alkylodioxy, amino, lower alkylamino,  
lower aminoalkyl, arylamino, nitro, and halosulfonyl; and

HetAr<sup>2</sup> is pyridinyl or pyrimidinyl optionally  
20 substituted with one or more radicals independently  
selected from lower alkylthio, lower alkylsulfonyl, lower  
alkylsulfinyl, halo, lower alkyl, lower heterocyclyl,  
lower alkoxy, lower aralkoxy, lower haloalkyl, amino,  
cyano, lower aralkyl, lower alkylamino, lower  
25 cycloalkylamino, lower arylamino, lower alkynylamino, and  
lower aralkylamino; or

a pharmaceutically-acceptable salt or tautomer  
thereof.

A class of compounds of particularly interest  
30 consists of these compounds of Formula I wherein

R<sup>1</sup> is selected from hydrido, methyl, ethyl,  
isopropyl, tert-butyl, isobutyl, trichloroethyl,  
pentafluoroethyl, heptafluoropropyl, difluoroethyl,  
difluoropropyl, dichloroethyl, dichloropropyl, vinyl,  
35 allyl, ethynyl, propargyl, morpholinyl, piperidinyl,  
piperazinyl, benzyl, phenylethyl, morpholinomethyl,

morpholinoethyl, pyrrolidinylmethyl, piperazinylmethyl, piperidinylmethyl, pyridinylmethyl, thienylmethyl, methoxymethyl, ethoxymethyl, amino, methylamino, dimethylamino, phenylamino, methylaminomethyl, dimethylaminomethyl, methylaminoethyl, dimethylaminoethyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclohexylmethyl, hydroxymethyl, hydroxyethyl, methylthio, and methylthiomethyl; and

$R^2$  is selected from hydrido, methyl, ethyl, propyl, isopropyl, tert-butyl, isobutyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, amino, N-methylamino, N,N-dimethylamino, ethynylamino, propargylamino, piperidinyl, piperazinyl, morpholinomethyl, pyrrolidinylmethyl, piperazinylmethyl, piperidinylmethyl, pyridinylmethyl, thienylmethyl, thiazolylmethyl, oxazolylmethyl, pyrimidinylmethyl, quinolylmethyl, isoquinolinylmethyl, imidazolylmethyl, benzimidazolylmethyl, furylmethyl, pyrazinylmethyl, aminocarbonylamino, methylaminocarbonylamino, dimethylaminocarbonylamino, ethylaminocarbonylamino, diethylaminocarbonylamino, methylsulfonylamino, ethylsulfonylamino, aminosulfonylamino, methylaminosulfonylamino, dimethylaminosulfonylamino, ethylaminosulfonylamino, and diethylaminosulfonylamino; and

$Ar^1$  is selected from phenyl, biphenyl, and naphthyl, wherein  $Ar^1$  is optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, aminosulfonyl, methyl, ethyl, isopropyl, tert-butyl, isobutyl, cyano, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl, methylcarbonylamino, trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl,

dichloromethyl, chloromethyl, allyl, vinyl, ethynyl, propargyl, methoxy, ethoxy, propyloxy, n-butoxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, aminomethyl, aminoethyl, N-methyl, N-phenylamino, phenylamino, diphenylamino, nitro, and chlorosulfonyl; and

HetAr<sup>2</sup> is selected from pyridinyl and pyrimidinyl, wherein HetAr<sup>2</sup> is optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, methyl, ethyl, isopropyl, tert-butyl, isobutyl, methoxyl, ethoxyl, phenoxyl, benzoxyl, phenethyl, trifluoromethyl, fluoromethyl, difluoromethyl, amino, benzylamino, propargylamino, cyclopropylamino, cyclobutylamino, cyclopentylamino, and cyano; or

a pharmaceutically-acceptable salt or tautomer thereof.

A class of compounds of specific interest consists of those compounds of Formula I wherein

R<sup>1</sup> is hydrido, methyl, ethyl, hydroxyethyl, propargyl, dimethylaminoethyl or morpholinoethyl; and

R<sup>2</sup> is selected from hydrido, methyl, ethyl, amino, aminocarbonylamino, methylaminocarbonylamino, methylsulfonylamino, aminosulfonylamino, and methylaminosulfonylamino; and

Ar<sup>1</sup> is phenyl optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, aminosulfonyl, methyl, ethyl, isopropyl, tert-butyl, isobutyl, cyano, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl, methylcarbonylamino, trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl, dichloromethyl, chloromethyl, methoxy, ethoxy, propyloxy, n-butoxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, aminomethyl, aminoethyl, N-methyl, N-

phenylamino, phenylamino, diphenylamino, nitro, and  
chlorosulfonyl; and

HetAr<sup>2</sup> is optionally substituted with one or more  
radicals independently selected from methylthio,  
5 methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo,  
methyl, ethyl, isopropyl, tert-butyl, isobutyl, methoxyl,  
ethoxyl, phenoxyl, benzoxyl, trifluoromethyl,  
fluoromethyl, difluoromethyl, amino, propargylamino, and  
cyano; or

10 a pharmaceutically-acceptable salt or a tautomer  
thereof.

A class of compounds of very specific interest  
consists of those compounds of Formula I wherein

R<sup>1</sup> is hydrido or methyl; and

15 R<sup>2</sup> is hydrido or methyl; and

Ar<sup>1</sup> is phenyl which is optionally substituted with  
one or more radicals independently selected fluoro,  
chloro, methyl, ethyl, trifluoromethyl, methoxy, ethoxy,  
dimethylamino, and nitro; and

20 HetAr<sup>2</sup> is optionally substituted with one or more  
radicals independently selected from methyl, chloro,  
fluoro, and trifluoromethyl; or

a pharmaceutically-acceptable salt or tautomer  
thereof.

25 A family of specific compounds of particular  
interest within Formula I consists of compounds, and  
tautomers and pharmaceutically-acceptable salts thereof,  
as follows:

4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine;

30 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine;

N-[4(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-  
yl]methanesulfonamide;

N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-  
N'-methylsulfamide;

35 [4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]urea;



- [4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl] sulfamide;  
4-(4-chlorophenyl)-1-methyl-3-(4-pyridinyl)-1H-pyrazol-5-amine;  
5 N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylurea;  
4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine;  
4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine;  
4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazole-1-ethanol;  
10 4-(4-fluorophenyl)-N,N-dimethyl-3-(4-pyridinyl)-1H-pyrazole-1-ethanamine;  
4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine;  
4-[4-(4-chlorophenyl)-1H-pyrazol-3-yl]pyridine;  
15 4-(4-phenyl-1H-pyrazol-5-yl)pyridine;  
1-methyl-4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine; and  
1-methyl-4-[2-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine.

- 20 The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH<sub>2</sub>-) radical. Where used, either alone or  
25 within other terms such as "haloalkyl", "alkylsulfonyl", "alkoxyalkyl", "hydroxyalkyl", "mercaptoalkyl", the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are  
30 "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl  
35 and the like. The term "alkenyl" embraces linear or branched radicals having at least one carbon-carbon



double bond of two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkenyl radicals are "lower alkenyl" radicals having two to about six carbon atoms. Examples of alkenyl radicals include ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The terms "alkenyl" and "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations. The term "alkynyl" embraces linear or branched radicals having at least one carbon-carbon triple bond of two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about six carbon atoms. Examples of alkynyl radicals include propargyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butenyl and 1-pentynyl. The term "cycloalkyl" embraces saturated carbocyclic radicals having three to about twelve carbon atoms. More preferred cycloalkyl radicals are "lower cycloalkyl" radicals having three to about eight carbon atoms. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term "cycloalkylalkylene" embraces alkyl radicals substituted with a cycloalkyl radical. More preferred cycloalkylalkylene radicals are "lower cycloalkylalkylene" which embrace lower alkyl radicals substituted with a lower cycloalkyl radical as defined above. Examples of such radicals include cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl. The term "cycloalkenyl" embraces partially unsaturated carbocyclic radicals having three to twelve carbon atoms and one or two double bonds but not necessarily conjugated ("cycloalkyldienyl"). More preferred cycloalkenyl radicals are "lower cycloalkenyl" radicals having four to about eight carbon atoms. Examples of such radicals include cyclobutenyl,

cyclopentenyl and cyclohexenyl. The term "cycloalkenylalkylene" embraces alkyl radicals substituted with a cycloalkenyl radical. More preferred cycloalkenylalkylene radicals are "lower

5 cycloalkenylalkylene" which embrace lower alkyl radicals substituted with a lower cycloalkenyl radical, as defined above. Examples of such radicals include cyclobutenylmethyl, cyclopentenylmethyl and cyclohexenylmethyl. The term "halo" means halogens such

10 as fluorine, chlorine, bromine or iodine. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl

15 radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. "Lower haloalkyl" embraces radicals having one to six

20 carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl,

25 difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms, any one of which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl

30 radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl. The terms "alkoxy" and "alkyloxy" embrace

35 linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More

preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and tert-butoxy. The term "alkoxyalkyl" embraces alkyl radicals having one or more alkoxy radicals attached to the alkyl radical to form, for example, monoalkoxyalkyl and dialkoxyalkyl radicals. The "alkoxy" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" radicals.

The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused. More preferred aryl are 6-12 membered aryl radicals. Examples of such radicals include phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. Aryl moieties may also be substituted at a substitutable position with one or more substituents selected independently from, for example, halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino, aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl, alkoxyalkyl, alkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, aralkoxy, hydroxyl, acyl, carboxy, aminocarbonyl, and aralkoxycarbonyl. The term "alkylldioxy" encompasses an alkylldioxy bridge, such as a methylenedioxy bridge, between two carbon ring atoms of an aryl moiety.

The term "heterocyclyl" embraces saturated, partially unsaturated and unsaturated heteroatom-containing ring-shaped radicals, which can also be called "heterocyclyl", "heterocycloalkenyl" and "heteroaryl" correspondingly, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated

heterocyclyl radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms (e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. morpholinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., thiazolidinyl, etc.). Examples of partially unsaturated heterocyclyl radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole. Heterocyclyl radicals may include a pentavalent nitrogen, such as in tetrazolium and pyridinium radicals. The term "heteroaryl" embraces unsaturated heterocyclyl radicals. Examples of heteroaryl radicals include unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.) tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclyl group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo[1,5-b]pyridazinyl, etc.), etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.; unsaturated 3 to 6-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc.; unsaturated 3- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. benzoxazolyl,

benzoxadiazolyl, etc.); unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl (e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like. The term "heteroaryl" also embraces radicals where heterocyclyl radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like. Said heterocyclyl group may have 1 to 3 substituents such as alkyl, hydroxyl, halo, alkoxy, oxo, amino and alkylamino. The term "heterocyclylalkylene" embraces heterocyclyl-substituted alkyl radicals. More preferred heterocyclylalkylene radicals are "lower heterocyclylalkylene" radicals having one to six carbon atoms and a heterocyclyl radical.

The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to about ten carbon atoms attached to a divalent sulfur atom. More preferred alkylthio radicals are "lower alkylthio" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthio radicals are methylthio, ethylthio, propylthio, butylthio and hexylthio. The term "alkylthioalkylene" embraces radicals containing an alkylthio radical attached through the divalent sulfur atom to an alkyl radical of one to about ten carbon atoms. More preferred alkylthioalkylene radicals are "lower alkylthioalkylene" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthioalkylene radicals include methylthiomethyl. The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to about ten carbon atoms, attached to a divalent -S(=O)- radical. More preferred alkylsulfinyl radicals

are "lower alkylsulfinyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylsulfinyl radicals include methylsulfinyl, ethylsulfinyl, butylsulfinyl and hexylsulfinyl. The term "sulfonyl", whether used alone or linked to other terms such as "alkylsulfonyl", or "halosulfonyl" denotes a divalent radical,  $-SO_2-$ . "Alkylsulfonyl" embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfonyl radicals include methylsulfonyl, ethylsulfonyl and propylsulfonyl. The "alkylsulfonyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkylsulfonyl radicals. The term "halosulfonyl" embraces halo radicals attached to a sulfonyl radical. Examples of such halosulfonyl radicals include chlorosulfonyl and bromosulfonyl. The terms "sulfamyl", "aminosulfonyl" and "sulfonamidyl" denote  $NH_2O_2S-$ .

The term "carbonyl", whether used alone or with other terms, such as "alkoxycarbonyl", denotes  $-(C=O)-$ . The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes  $-CO_2H$ . The term "carboxyalkyl" embraces alkyl radicals substituted with a carboxy radical. More preferred are "lower carboxyalkyl" radicals which embrace carboxy-substituted lower alkyl radicals, as defined above. Examples of such lower carboxyalkyl radicals include carboxymethyl, carboxyethyl and carboxypropyl. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. More preferred are "lower alkoxycarbonyl" radicals with alkyl portions having one to six carbons. Examples of such lower alkoxycarbonyl (ester) radicals include methoxycarbonyl, ethoxycarbonyl,



propoxycarbonyl, butoxycarbonyl and hexyloxycarbonyl. The term "alkoxycarbonylalkylene" embraces alkyl radicals substituted with an alkoxycarbonyl radical as defined above. More preferred are "lower alkoxycarbonylalkylene" radicals with alkyl portions having one to six carbons. Examples of such lower alkoxycarbonylalkylene radicals include methoxycarbonylmethylene, ethoxycarbonylmethylene, methoxycarbonylethylene and ethoxycarbonylethylene. The term "alkylcarbonyl", includes radicals having alkyl radicals attached to a carbonyl radical. Examples of such radicals include methylcarbonyl, ethylcarbonyl, propylcarbonyl, butylcarbonyl, and pentylcarbonyl. The term "aralkyl" embraces aryl-substituted alkyl radicals. Preferred aralkyl radicals are "lower aralkyl", having lower alkyl groups substituted with one or more aryl groups. Examples of such groups include benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, and diphenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The terms benzyl and phenylmethyl are interchangeable. The term "heterocyclylalkylene" embraces saturated, partially unsaturated and unsaturated heterocyclyl-substituted alkyl radicals such as pyrrolidinylmethyl, pyridylmethyl, quinolylmethyl, thienylmethyl, furylethyl, and quinolylethyl. The heteroaryl in heteroaralkyl (unsaturated heterocyclyl-substituted alkyl radicals) may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The term "aryloxy" embraces aryl radicals attached through an oxygen atom to other radicals. The term "aralkoxy" embraces aralkyl radicals attached through an oxygen atom to other radicals.

The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. More preferred are "lower aminoalkyl" radicals. Examples of such radicals include aminomethyl, aminoethyl, and the like. The term



"alkylamino" denotes amino groups which are substituted with one or two alkyl radicals. Preferred are "lower alkylamino" radicals having alkyl portions having one to six carbon atoms. Suitable lower alkylamino may be

5 monosubstituted N-alkylamino or disubstituted N,N-alkylamino, such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like. The term "arylamino" denotes amino groups which are substituted with one or two aryl radicals, such as N-phenylamino.

10 The "arylamino" radicals may be further substituted on the aryl ring portion of the radical. The term "aminocarbonyl" denotes an amide group of the formula -C(=O)NH. The term "alkylaminocarbonyl" denotes an aminocarbonyl group which has been substituted with one

15 or two alkyl radicals on the amino nitrogen atom. Preferred are "N-alkylaminocarbonyl" and "N,N-dialkylaminocarbonyl" radicals. More preferred are "lower N-alkylaminocarbonyl" and "lower N,N-dialkylaminocarbonyl" radicals with lower alkyl portions

20 as defined above. The term "aminocarbonylamino" embraces radicals having one or more aminocarbonyl radicals attached to an amino radical. The term "alkylaminocarbonylamino" embraces radicals having one or more alkyl radicals attached to an aminocarbonylamino

25 radical. Preferred are "lower alkylaminocarbonylamino" radicals with lower alkyl portions as defined above. The term "alkylcarbonylamino" embraces amino groups which are substituted with one or more alkylcarbonyl radicals. More preferred alkylcarbonylamino radicals are "lower

30 alkylcarbonylamino" having lower alkylcarbonyl radicals as defined above attached to amino radicals. The term "alkylaminoalkylene" embraces radicals having one or more alkyl radicals attached to an aminoalkyl radical. The term "alkylsulfonylamino" embraces radicals having one or


35 more alkylsulfonyl radicals attached to an amino radical. Preferred are "lower alkylsulfonylamino" radicals with

lower alkyl portions as defined above. The term "aminosulfonylamino" embraces radicals having one or more aminosulfonyl radicals attached to an amino radical. The term "alkylaminosulfonylamino" embraces radicals having  
5 one or more alkyl radicals attached to an aminosulfonylamino radical. Preferred are "lower alkylaminosulfonylamino" radicals with lower alkyl portions as defined above.

The additional terms used to describe the  
10 substituents of the pyrazole ring and not specifically defined herein are defined in a similar manner to that illustrated in the above definitions. As above, more preferred substituents are those containing "lower" radicals. Unless otherwise defined to contrary, the term  
15 "lower" as used in this application means that each alkyl radical of a pyrazole ring substituent comprising one or more alkyl radicals has one to about six carbon atoms; each alkenyl radical of a pyrazole ring substituent comprising one or more alkenyl radicals has two to about  
20 six carbon atoms; each alkynyl radical of a pyrazole ring substituent comprising one or more alkynyl radicals has two to about six carbon atoms; each cycloalkyl or cycloalkenyl radical of a pyrazole ring substituent comprising one or more cycloalkyl and/or cycloalkenyl  
25 radicals is a 3 to 8 membered ring cycloalkyl or cycloalkenyl radical, respectively; each aryl radical of a pyrazole ring substituent comprising one or more aryl radicals is a monocyclic aryl radical; and each heterocyclyl radical of a pyrazole ring substituent  
30 comprising one or more heterocyclyl radicals is a 4-8 membered ring heterocyclyl.

The present invention comprises the tautomeric forms of compounds of Formula I. As illustrated below, the pyrazoles of Formula I and I' are magnetically and  
35 structurally equivalent because of the prototropic tautomeric nature of the hydrogen:

23




The present invention also comprises compounds of Formula I having one or more asymmetric carbons. It is known to those skilled in the art that those pyrazoles of the present invention having asymmetric carbon atoms may exist in diastereomeric, racemic, or optically active forms. All of these forms are contemplated within the scope of this invention. More specifically, the present invention includes enantiomers, diastereomers, racemic mixtures, and other mixtures thereof.

The present invention comprises a pharmaceutical composition for the treatment of a TNF mediated disorder, a p38 kinase mediated disorder, inflammation, and/or arthritis, comprising a therapeutically-effective amount of a compound of Formula I, or a therapeutically-acceptable salt or tautomer thereof, in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a therapeutic method of treating a TNF mediated disorder, a p38 kinase mediated disorder, inflammation and/or arthritis in a subject, the method comprising treating a subject having or susceptible to such disorder or condition with a therapeutically-effective amount of a compound of Formula I

24



wherein

$R^1$  is selected from hydrido, alkyl, cycloalkyl,  
 5 alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene,  
 haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl,  
 mercaptoalkyl, alkylthioalkylene, amino, alkylamino,  
 arylamino, aminoalkyl, alkylaminoalkylene,  
 heterocyclalkylene, aminocarbonylalkylene, and  
 10 alkylaminocarbonylalkylene; and

$R^2$  is selected from hydrido, alkyl, alkenyl, alkynyl,  
 heterocyclyl, haloalkyl, heterocyclalkyl, amino,  
 alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy,  
 alkoxycarbonyl, carboxyalkyl, aminocarbonylamino,  
 15 alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl,  
 alkylsulfonylamino, aminosulfonylamino,  
 alkylaminosulfonylamino, and alkynylamino; wherein the  
 heterocyclyl and heterocyclalkyl groups are optionally  
 substituted with one or more radicals independently  
 20 selected from alkylthio, alkylsulfonyl, alkylsulfinyl,  
 halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl,  
 haloalkyl, amino, cyano, and hydroxy; and

$Ar^1$  is aryl optionally substituted with one or more  
 radicals independently selected from halo, alkyl,  
 25 alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy,  
 alkylthio, alkylsulfinyl, alkylsulfonyl, amino,  
 aminocarbonyl, cyano, alkoxycarbonyl, formyl,  
 aminosulfonyl, alkylamino, nitro, arylamino,  
 alkylcarbonylamino, halosulfonyl, aminoalkyl, and  
 30 haloalkyl; and

HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

Also included in the family of compounds of Formula I are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclyl, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, stearic, salicylic, *p*-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, cyclohexylaminosulfonic, algenic,  $\beta$ -hydroxybutyric, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts and organic


salts. More preferred metallic salts include, but are not limited to appropriate alkali metal (group Ia) salts, alkaline earth metal (group IIa) salts and other physiological acceptable metals. Such salts can be made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc. Preferred organic salts can be made from tertiary amines and quaternary ammonium salts, including in part, tromethamine, diethylamine, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

#### General Synthetic Procedures

The compounds of the invention can be synthesized according to the following procedures of Schemes I - VI wherein the  $R^1$  -  $R^3$  substituents and  $Ar^1$ ,  $HetAr^2$  are as defined for Formula I, above, except where further noted.

#### Scheme I


27



Scheme I shows the three step preparation of the pyrazole 5 of the present invention. In step 1, the reaction of arylmethyl derived ketone 1 with pyridine derived aldehyde 2 either in a solvent such as benzene or toluene in the presence of a base such as pyridine or in a mixture of acids such as acetic acid and hydrogen bromide gives the  $\alpha,\beta$ -unsaturated ketone 3. In step 2, in the presence of base such as sodium hydroxide,  $\alpha,\beta$ -unsaturated ketone 3 is converted to the corresponding epoxide 4 by the treatment with hydrogen peroxide solution at room temperature. In step 3, epoxide 4 is condensed with hydrazine in a suitable solvent such as ethanol at temperature ranging up to the boiling point to form pyrazole 5. Alternatively, pyrazole 5 can be prepared by treatment of 3 with tosyl hydrazide in the presence of an acid such as acetic acid at reflux.




28



Scheme II shows the synthesis of pyrazole 12 containing a heteroaromatic ring by three routes. In Route 1, ketone 6 is condensed with hydrazine 7 to give substituted hydrazine 9, which is then reacted with acyl halide or anhydride 10 at low temperature to provide acyl hydrazone 11. Upon heating at temperature up to 200 °C, hydrazone 11 is converted to pyrazole 12. In Route 2, acyl hydrazone 11 is formed directly by reaction of ketone 6 with acyl hydrazide 8 at room temperature. Acyl hydrazide 8 may be formed by reaction of hydrazine with a carboxylic acid ester. Heating 11 as above then provides pyrazole 12. In Route 3, ketone 6 is treated with acyl hydrazide 8 at from room temperature to ~200 °C to give pyrazole 12 directly. Alternatively, this condensation may be carried out in an acidic solvent, such as acetic acid, or in a solvent containing acetic acid.

Scheme III




- Cyanoketone **13** may be synthesized according to the procedure described by I. Lantos et al in J. Org. Chem., volume 53, pp. 4223-4227 (1988) for the synthesis of the p-fluoro compound (X = p-F). This procedure, which is incorporated herein by reference, can be used to synthesize cyanoketones such as **13** wherein X is selected from, for example, halogen, alkyl and alkoxy.
- Cyanoketones such as **13** may be converted to pyrazoles **14** by reaction with a hydrazine in a suitable solvent, such as benzene or toluene. A catalyst such as acetic acid may be employed. When hydrazine itself is employed, the ring nitrogen atoms of the pyrazole thus obtained bear no substituent except hydrogen on one of the ring nitrogen atoms. When a substituted hydrazine, such as methylhydrazine is employed, the product pyrazole **14** bears a substituent on the ring nitrogen atom adjacent to the aminated ring carbon atom, as shown in Scheme 1. The resultant aminopyrazole **14** may be acylated or sulfonylated to form substituted aminopyrazole **15** by treatment with a suitably activated carboxylic or sulfonic acid in a suitable solvent such as pyridine. Examples of a suitably activated carboxylic acid include

acetic anhydride or benzoyl chloride. Examples of a suitably activated sulfonic acid include methanesulfonyl chloride, p-toluenesulfonyl chloride or sulfamyl chloride.

5


## Scheme IV



Scheme IV illustrates the synthesis of 3-pyridyl-4-aryl-pyrazoles of the present invention. Benzoate 16 is first reacted with pyridine 17 in the presence of a base, such as an alkali metal alkoxide (preferably sodium methoxide), in a suitable solvent, such as tetrahydrofuran. Subsequent treatment with an acid, preferably a mineral acid such as hydrochloric acid, yields the desoxybenzoin 18. Desoxybenzoin 18 is then converted to ketone 19 by treatment with an excess of dimethylformamide dimethyl acetal. Ketone 19 is then reacted with hydrazine in a suitable solvent such as ethanol to yield a mixture of pyrazoles 20 and 21. In Scheme IV, R<sup>4</sup> represents one or more radicals independently selected from the optional substituents previously defined for Ar<sup>1</sup>; and R<sup>5</sup> represents one or more radicals independently selected from the optional substituents previously defined for HetAr<sup>2</sup>.


The 3-pyrimidinyl-4-aryl-pyrazoles of the present invention can be synthesized in the manner of Scheme IV by replacing pyridine 17 with the corresponding pyrimidine.

## Scheme V



In Scheme V, hydroxyalkyl pyrazoles 22 and 23 are converted to sulfonate derivatives by reaction with an alkyl- or arylsulfonyl halide. These sulfonates are then reacted with ammonia or primary amines or secondary amines to give the corresponding 1-amino-pyrazoles 24 and 25, respectively. In Scheme V, n is 1, 2, 3, 4 or 5; R<sup>4</sup> and R<sup>5</sup> are as defined in Scheme IV; R<sup>6</sup> and R<sup>7</sup> are independently selected, for example, from hydrogen, alkyl and aryl, or together with the nitrogen atom to which they are attached form a 4-8 membered ring that may contain one or more additional heteroatoms selected from oxygen, nitrogen or sulfur.

Scheme VI



5        Scheme VI is similar to Scheme IV except that  
 desoxybenzoin 18 is first reacted with hydrazine in a  
 suitable solvent such as ethanol to yield hydrazine 26.  
 Hydrazine 26 is then converted to pyrazole 20 (rather  
 than a mixture of pyrazoles 20 and 21 as in Scheme IV) by  
 10 treatment with an excess of dimethylformamide dimethyl  
 acetal. In Scheme VI, R<sup>4</sup> and R<sup>5</sup> are as defined in Scheme  
 V.

15        The following examples contain detailed descriptions  
 of the methods of preparation of compounds of Formula I.  
 These detailed descriptions fall within the scope, and  
 serve to exemplify, the above described General Synthetic  
 Procedures which form part of the invention. These  
 detailed descriptions are presented for illustrative  
 purposes only and are not intended as a restriction on


the scope of the invention. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated. All compounds showed NMR spectra consistent with their assigned structures. In some cases, the  
5 assigned structures were confirmed by nuclear Overhauser effect (NOE) experiments.

The following abbreviations are used:

HCl - hydrochloric acid  
MgSO<sub>4</sub> - magnesium sulfate  
10 Na<sub>2</sub>SO<sub>4</sub> - sodium sulfate  
NaIO<sub>4</sub> - sodium periodate  
NaHSO<sub>3</sub> - sodium bisulfite  
NaOH - sodium hydroxide  
KOH - potassium hydroxide  
15 P<sub>2</sub>O<sub>5</sub> - phosphorus pentoxide  
MeOH - methanol  
EtOH - ethanol  
HOAc (or AcOH) - acetic acid  
EtOAc - ethyl acetate  
20 H<sub>2</sub>O - water  
H<sub>2</sub>O<sub>2</sub> - hydrogen peroxide  
CH<sub>2</sub>Cl<sub>2</sub> - methylene chloride  
NaOMe - sodium methoxide  
h - hour  
25 hr - hour  
min - minutes  
THF - tetrahydrofuran  
TLC - thin layer chromatography  
DSC - differential scanning calorimetry  
30 b.p. - boiling point  
m.p. - melting point  
eq - equivalent



## EXAMPLE 1



## 4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine

5

Step 1: Preparation of 3-phenyl-4-(4-pyridyl)-3-butene-2-one

3-Phenyl-4-(4-pyridyl)-3-butene-2-one was prepared by the method of Reichert and Lechner, Arzneim.-Forsch. **15**, 36  
10 (1965), which is incorporated by reference herein.

Step 2: Preparation of 3-phenyl-4-(4-pyridyl)-3,4-epoxy-2-butanone

To a stirred solution of 3-phenyl-4-(4-pyridyl)-3-butene-2-one (step 1) (500 mg, 2.24 mmol) in methanol (10  
15 ml) at room temperature was added an aqueous solution (9 ml) of sodium hydroxide (100 mg, 2.24 mmol) and hydrogen peroxide (0.5 ml of 30% aqueous solution, 4.4 mmol). After stirring for 2 hours, sodium chloride was added and  
20 the reaction was extracted with ethyl acetate. The combined organic layers were dried over magnesium sulfate, filtered, and concentrated in vacuo to provide the crude 3-phenyl-4-(4-pyridyl)-3,4-epoxy-2-butanone (385 mg, 65%) as an oil. This was used in the next step  
25 without further purification.


Step 3: Preparation of 4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine

A solution of 3-phenyl-4-(4-pyridyl)-3,4-epoxy-2-butanone (step 2) (350 mg, 1.46 mmol) and anhydrous  
30 hydrazine (0.7 ml, 20 mmol) in ethanol (3 ml) was heated at reflux for 4 hours. The reaction was cooled, and the

solvent was evaporated to dryness. The resulting residue was purified by chromatography (silica gel, 1:1 acetone/hexane) to give the desired product as a crystalline solid, which was recrystallized from ethyl acetate and hexane to give pure 4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine (145 mg, 42%): m. p. 164-165°C. Anal. Calc'd for  $C_{15}H_{13}N_3$  (235.29): C, 76.57; H, 5.57; N, 17.86. Found: C, 76.49; H, 5.45; N, 17.70.

The compounds of Examples 2 through 8 were synthesized in accordance with the chemistry described above (particularly in Scheme III) by selection of the corresponding starting reagents:

#### EXAMPLE 2



4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine


The cyanoketone 1 of Scheme III wherein X is p-fluoro was synthesized according to the procedure of I. Lantos et al., J. Org. Chem., **53**, 4223-4227 (1988), which is incorporated herein by reference. A solution of the cyanoketone (10 g, 41 mmol), hydrazine hydrate (2.5 ml) and acetic acid (5.2 ml) in benzene (100 ml) was refluxed for 4 hours. The reaction was cooled and extracted with 3N HCl. The combined acid extracts were basified to pH 10 using concentrated ammonium hydroxide with cooling. The basic aqueous layer was extracted with methylene chloride and the combined organic extracts were dried over magnesium sulfate.

The drying agent was filtered and the filtrate concentrated in vacuo to give the crude 4-(4-

fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine which was purified by recrystallization from ethyl acetate and hexane. Purified 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine had m.p. 178-180°C (capillary).

5           Anal. Calc'd for  $C_{14}H_{11}N_4F + 0.25 H_2O$ : C, 64.99; H, 4.48; N, 21.65. Found: C, 64.99; H, 4.48; N, 21.54.

### EXAMPLE 3



10


N-[4(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide

A solution of 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine prepared as set forth in Example 2 (200 mg, 0.77 mmol) and methanesulfonyl chloride (90 mg) in  
15   pyridine (5 ml) was stirred at room temperature overnight. The pyridine was removed in vacuo and the residue was treated with methylene chloride and water. The resultant precipitate was filtered to give N-[4(4-  
20   fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide. Additional N-[4(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide was contained in the methylene chloride layer. The methylene chloride was stripped in vacuo and the residue purified  
25   by chromatography on silica gel using mixtures of ethyl acetate and methanol as eluents. The purified N-[4(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide had m.p. 168-170°C.

30           Anal. Calc'd for  $C_{15}H_{13}N_4SO_2F + 0.25 H_2O$ : C, 53.48; H, 4.04; N, 16.63. Found: C, 53.41; H, 3.78; N, 16.52.

### EXAMPLE 4

38



N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylsulfamide

5       Methyl sulfamyl chloride was synthesized by refluxing a solution of methylsulfamic acid (1.0 g) in phosphorus oxychloride (10 mL) for 6 hours. The excess phosphorus oxychloride was removed in vacuo and the residual oil was used for the synthesis of the product


10       without further treatment. A solution of 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine prepared as set forth in Example 2 (200 mg, 0.77 mmol) and approximately 1 mmol of the above oil in pyridine (5 ml) was stirred at room temperature for 2 hours. The

15       reaction was stripped in vacuo and the residue purified by chromatography on silica gel using ethyl acetate and mixtures of ethyl acetate and methanol as eluents to give N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylsulfamide as a crystalline solid, m. p. 194-

20       195°C.

Anal. Calc'd for  $C_{15}H_{14}N_5SO_2F + 1.0 H_2O$ : C, 49.31; H, 4.41; N, 19.17. Found: C, 49.13; H, 3.97; N, 19.01.

## EXAMPLE 5



[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]urea


5        A suspension of 4-(4-fluorophenyl)-5-(4-pyridinyl)-  
1H-pyrazol-3-amine prepared as set forth in Example 2  
(200 mg, 0.77 mmol) in a solution of di-tert-butyl  
carbonate (185 mg, 0.9 mmol) and 4-dimethylaminopyridine  
(DMAP) (10 mg) in methylene chloride (10 ml) was stirred  
10    at room temperature for 20 minutes, during which time,  
the suspended material dissolved. N-Propylamine (50 mg)  
was added and stirring was continued at room temperature  
for 1 hour. The reaction was then refluxed for 15  
minutes, cooled and stripped in vacuo. Treatment with  
15    ethyl acetate and hexane resulted in the deposition of  
crystals of the tert-butoxycarbonyl derivative, m.p. 183-  
184°C.

Anal. Calc'd for  $C_{19}H_{19}N_4O_2F$ : C, 64.40; H, 5.40; N,  
15.81. Found: C, 64.66; H, 5.63; N, 15.63.

20        A solution of the tert-butoxycarbonyl derivative  
above (100 mg, 0.3 mmol) in tetrahydrofuran was treated  
with ammonia at 80°C in a pressure bottle for 12 hours.  
The reaction was stripped in vacuo and the residue was  
purified by chromatography on silica gel eluting with  
25    mixtures of ethyl acetate and methanol. The purified [4-  
(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]urea  
thus obtained had m.p. 224-225°C.

Anal. Calc'd for  $C_{15}H_{12}N_5O$ : C, 60.60; H, 4.07; N,  
23.56. Found: C, 60.21; H, 4.11; N, 23.30.

## EXAMPLE 6




5 [4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]sulfamide

Sulfamyl chloride was synthesized from chlorosulfonyl isocyanate according to the procedure described by R. Graf in Chemische Berichte, p. 509 (1959), which is incorporated herein by reference. A solution of 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine prepared as set forth in Example 2 (200 mg, 0.77 mmol), sulfamyl chloride (100 mg, 0.8 mmol) and triethylamine (200 mg, 2 mmol) in benzene (5 ml) and acetonitrile (5 ml) was stirred at room temperature for 2 hours. The reaction was stripped in vacuo and residue was treated with water and basified to pH 7 with ammonium hydroxide. The resultant precipitate was purified by chromatography on silica gel using mixtures of ethyl acetate and methanol as eluents. The purified [4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]sulfamide thus obtained had m.p. 201-202°C.

Anal. Calc'd for  $C_{14}H_{12}N_5SO_2F$ : C, 50.44; H, 3.63; N, 21.01. Found: C, 50.43; H, 3.45; N, 20.89.


41

**EXAMPLE 7**

5           4-(4-chlorophenyl)-1-methyl-3-(4-pyridinyl)-1H-pyrazol-5-amine

A solution of cyanoketone 1 of Scheme III wherein X is p-chloro (1.5 g, 5.19 mmol), methylhydrazine (0.35 ml) and acetic acid (0.75 ml) in benzene (15 ml) was refluxed  
10 for 3.5 hours. The reaction was cooled and extracted with 3N HCl. The aqueous layer was concentrated on the rotary evaporator and then basified with ammonium hydroxide. The resultant precipitate was recrystallized from methanol to give pure 4-(4-chlorophenyl)-1-methyl-3-(4-pyridinyl)-1H-pyrazol-5-amine, m.p. 257-258°C.  
15

Anal. Calc'd for  $C_{15}H_{13}N_4Cl$ : C, 63.27; H, 4.60; N, 19.68. Found: C, 62.93; H, 4.45; N, 19.64.

**EXAMPLE 8**

20

N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylurea




A solution of 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine prepared as set forth in Example 2 (100 mg, 0.38 mmol), methyl isocyanate (22 mg, 0.39 mmol) and 4-dimethylaminopyridine (2.5 mg) in methylene chloride (10 ml) was stirred at room temperature for 30 minutes. The reaction was stripped in vacuo. The residue was triturated with hexane and the solid filtered to give pure N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylurea, m. p. 212-213°C.

Anal. Calc'd for  $C_{16}H_{14}N_5FO$ : C, 61.73; H, 4.53; N, 22.50. Found: C, 61.63; H, 4.55; N, 22.47.

The compounds of Examples 9 through 11 were synthesized in accordance with the chemistry described above (particularly in Scheme IV) by selection of the corresponding starting reagents:

#### EXAMPLE 9



4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine

##### Step 1


Methyl isonicotinate (13.7g, 0.1mole) and ethyl 4-fluorophenylacetate (18.2g, 0.1mole) were mixed together, then sodium methoxide (8.1g, 0.15mole) was added. The mixture was heated to 60-70°C for 24 hours while nitrogen was blown through the flask to eliminate methanol. Concentrated hydrochloric acid (50 mL) then was added and the reaction mixture was refluxed for 3 hours. After

addition of water (30 mL), the reaction mixture was extracted with chloroform, and the water phase was neutralized to pH 6 - 7 with aqueous sodium hydroxide (1M). The precipitate formed was collected by  
5 filtration, washed with water and dried under vacuum to give 10 g of 2-(4-fluorophenyl)-1-(4'-pyridyl)-ethan-1-one (yield: 46%). <sup>1</sup>H NMR: consistent with the assigned structure and/or its tautomer.

10 Step 2


2-(4-fluorophenyl)-1-(4'-pyridyl)-ethan-1-one prepared above (1 g) was dissolved in 50 mL tetrahydrofuran and N,N-dimethylformamide dimethyl acetal (5 mL) was added. The mixture was stirred at room  
15 temperature for 2 days. After evaporating the solvent, the solid obtained was washed with hexane and 1 g of the vinyl amine was obtained. This vinyl amine (0.5 g) was dissolved in ethanol (15 mL) and hydrazine hydrate (5 mL) was added. The mixture was stirred at 0°C for 2 hours and  
20 then evaporated to dryness. After recrystallization from methanol/water, 400 mg of 4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine was obtained in 91% yield. MS, 240(M+1); <sup>1</sup>H NMR: consistent with the assigned structure; Anal. Calc'd for C<sub>14</sub>H<sub>10</sub>FN<sub>3</sub>.0.2H<sub>2</sub>O: C, 69.24; H, 4.32; N,  
25 17.30. Found: C, 69.54; H, 4.06; N, 17.43.

**EXAMPLE 10**



4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine

When methylhydrazine was substituted for hydrazine hydrate in Step 2 of Example 9, 4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine (the N-methyl derivative corresponding to the compound of Example 9) was obtained. Purification by recrystallization from toluene and hexane give the pure 4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine in 57% yield. MS m/z: 254 (M+1). <sup>1</sup>H NMR: consistent with the assigned structure. Anal. calc'd for C<sub>15</sub>H<sub>12</sub>FN<sub>3</sub>: C, 71.13; H, 4.78; N, 16.69. Found: C, 70.99; H, 4.68; N, 16.65.


**EXAMPLE 11**

4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazole-1-ethanol  
and  
4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazole-1-ethanol

The procedure set forth in Example 9 was followed except that 2-hydroxyethyl hydrazine was substituted for hydrazine hydrate. 4-(4-Fluorophenyl)-3-(4-pyridinyl)-1H-pyrazole-1-ethanol and 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazole-1-ethanol were obtained as a mixture by recrystallization from toluene and hexane in 67% yield. <sup>1</sup>H NMR: consistent with the assigned structure. Mass spectrum, m/z: 284 (M+1). Anal. calc'd for C<sub>16</sub>H<sub>14</sub>FN<sub>3</sub>O: C, 67.83; H, 4.98; N, 14.83. Found: C, 67.86; H, 5.04; N, 14.85.

The compounds of Examples 12 and 13 were synthesized in accordance with the chemistry described above (particularly in Scheme V) by selection of the corresponding starting reagents:

5

**EXAMPLE 12**

4-(4-fluorophenyl)-N,N-dimethyl-3-(4-pyridinyl)-1H-pyrazole-1-ethanamine

10

and


4-(4-fluorophenyl)-N,N-dimethyl-5-(4-pyridinyl)-1H-pyrazole-1-ethanamine

15

4-(4-Fluorophenyl)-3-(4-pyridinyl)-1H-pyrazole-1-ethanol (or 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazole-1-ethanol) prepared as set forth in Example 11 (1.36 g) was dissolved in 30 mL pyridine and cooled to 0°C, whereupon methanesulfonyl chloride (0.6 mL) was added. After stirring at 0°C for 12 hours, about 20 g of ice was added, and the mixture was extracted with toluene (300 ml). After evaporation, the residue was used directly without further purification. 0.7 g of the above obtained compound was dissolved in methanol (25 mL), and dimethylamine/THF solution (4M, 2 mL) was added. The reaction mixture was refluxed for 12 hours, then evaporated to dryness. The residue was purified by chromatography (methanol/dichloromethane 1:10). A

25

mixture (0.59 g) of 4-(4-fluorophenyl)-N,N-dimethyl-3-(4-pyridinyl)-1H-pyrazole-1-ethanamine and 4-(4-fluorophenyl)-N,N-dimethyl-5-(4-pyridinyl)-1H-pyrazole-1-ethanamine were obtained. <sup>1</sup>H NMR: consistent with the assigned structure. Mass spectrum, m/z: 311 (M+1). Anal. calc'd C<sub>18</sub>H<sub>19</sub>N<sub>4</sub>F.0.55H<sub>2</sub>O: C, 67.50; H, 6.33; N, 17.49. Found: C, 67.21; H, 6.46; N, 17.14.

**EXAMPLE 13**

4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine


and

4-[2-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine

The procedure set forth in Example 11 was followed, except that morpholine was substituted for dimethylamine, to produce a mixture of 4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine and 4-[2-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine. Mass spectrum, m/z: 353 (M+1). Anal. calc'd for C<sub>20</sub>H<sub>21</sub>N<sub>4</sub>OF + 0.5H<sub>2</sub>O: C, 66.47; H, 6.14; N, 15.50. Found: C, 66.57; H, 6.27; N, 15.14.

The compound of Example 14 was synthesized in accordance with the chemistry described above (particularly in Scheme VI) by selection of the corresponding starting reagents:

5

**EXAMPLE 14**

4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine


10

2-(4-fluorophenyl)-1-(4'-pyridyl)-ethan-1-one prepared as set forth in step 1 of Example 9 (0.5 g, 0.00232 moles) was mixed with of 98% methyl hydrazine (0.2 g, 0.00462 moles) in 10 mL of ethanol containing 0.1 mL of acetic acid in a 50 mL Erlenmeyer flask. After gentle boiling (30 minutes on a steam bath) a small sample was evacuated at high vacuum and examined by NMR to confirm completion of hydrazone formation. The reaction mixture was concentrated to a pasty mass and 3.6 mL of DMF dimethylacetal (0.027 moles) was then added and heated to 80°C for 30 minutes, at which point a clear yellow viscous solution was obtained. The reaction was checked for completion (TLC or NMR) and concentrated and taken up in 20 mL of chloroform. After washing with water (10 mL), the organic layer was extracted with 15 mL of 10 % HCl. The water layer was then treated with 0.5 g of activated charcoal at 70°C for 10 minutes, filtered through celite, neutralized cautiously to pH 7-8 with

25

vigorous stirring and cooling. The fine off-white precipitate was filtered and dried. NMR was found to be in agreement with the proposed structure. The precipitate, 4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine, obtained in quantitative yield was filtered, washed with ether and dried. Yield: 0.45 g (77%). Mass spectrum, m/z: 254. Anal. calc'd: C, 62.18; H, 4.52; N, 14.50. Found: C, 62.39; H, 4.07; N, 14.24.

10

**EXAMPLE 15**


4-[4-(4-chlorophenyl)-1H-pyrazol-3-yl]pyridine

15

4-[4-(4-chlorophenyl)-1H-pyrazol-3-yl]pyridine was prepared according to the procedure set forth in Example 9 except that ethyl 4-chlorophenylacetate was substituted for ethyl 4-fluorophenylacetate; m.p. 204-207°C.

20

Anal. Calc'd: C, 65.76; H, 3.94; N, 16.43. Found: C, 65.44; H, 3.78; N, 16.04.

**EXAMPLE 16**


25

4-(4-phenyl-1H-pyrazol-5-yl)pyridine



4-(4-phenyl-1H-pyrazol-5-yl)pyridine can be prepared in accordance with the procedure set forth in Example 9 by substituting ethylphenylacetate for ethyl 4-fluorophenylacetate.

5

**EXAMPLE 17**

10 1-methyl-4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine

and

1-methyl-4-[2-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine

15

This compound can be prepared using the procedure set forth for the synthesis of the compound of Example 11 by substituting 4-hydrazino-N-methylpiperidine for hydroxyethyl hydrazine. 4-Hydrazino-N-methylpiperidine is synthesized as disclosed in Ebnoether et al, Helv. Chim. Acta (1959) 42, 533, 541, 560. The resulting mixture is separated into the respective pure title compounds by chromatography on silica gel, eluting with methanol/dichloromethane (1:10), or other suitable solvent system.

20

25

BIOLOGICAL EVALUATION**p38 Kinase Assay**5    Cloning of human p38a:

The coding region of the human p38a cDNA was obtained by PCR-amplification from RNA isolated from the human monocyte cell line THP.1. First strand cDNA was synthesized from total RNA as follows: 2  $\mu$ g of RNA was  
10 annealed to 100 ng of random hexamer primers in a 10  $\mu$ l reaction by heating to 70 °C for 10 minutes followed by 2 minutes on ice. cDNA was then synthesized by adding 1  $\mu$ l of RNasin (Promega, Madison WI), 2  $\mu$ l of 50 mM dNTP's, 4  $\mu$ l of 5X buffer, 2  $\mu$ l of 100 mM DTT and 1  $\mu$ l (200 U) of  
15 Superscript II <sup>TM</sup> AMV reverse transcriptase. Random primer, dNTP's and Superscript <sup>TM</sup> reagents were all purchased from Life-Technologies, Gaithersburg, MA. The reaction was incubated at 42 °C for 1 hour.

Amplification of p38 cDNA was performed by aliquoting 5  
20  $\mu$ l of the reverse transcriptase reaction into a 100  $\mu$ l PCR reaction containing the following: 80  $\mu$ l dH<sub>2</sub>O, 2  $\mu$ l 50 mM dNTP's, 1  $\mu$ l each of forward and reverse primers (50 pmol/ $\mu$ l), 10  $\mu$ l of 10X buffer and 1  $\mu$ l Expand <sup>TM</sup> polymerase (Boehringer Mannheim). The PCR primers  
25 incorporated Bam HI sites onto the 5' and 3' end of the amplified fragment, and were purchased from Genosys. The sequences of the forward and reverse primers were 5'-GATCGAGGATTCATGTCTCAGGAGAGGCCCA-3' and  
5'-GATCGAGGATTCTCAGGACTCCATCTCTTC-3' respectively. The  
30 PCR amplification was carried out in a DNA Thermal Cycler (Perkin Elmer) by repeating 30 cycles of 94 °C for 1 minute, 60 °C for 1 minute and 68 °C for 2 minutes. After amplification, excess primers and unincorporated dNTP's were removed from the amplified fragment with a  
35 Wizard <sup>TM</sup> PCR prep (Promega) and digested with Bam HI (New England Biolabs). The Bam HI digested fragment was

ligated into BamHI digested pGEX 2T plasmid DNA (PharmaciaBiotech) using T-4 DNA ligase (New England Biolabs) as described by T. Maniatis, *Molecular Cloning: A Laboratory Manual*, 2nd ed. (1989). The ligation reaction was transformed into chemically competent *E. coli* DH10B cells purchased from Life-Technologies following the manufacturer's instructions. Plasmid DNA was isolated from the resulting bacterial colonies using a Promega Wizard<sup>TM</sup> miniprep kit. Plasmids containing the appropriate Bam HI fragment were sequenced in a DNA Thermal Cycler (Perkin Elmer) with Prism<sup>TM</sup> (Applied Biosystems Inc.). cDNA clones were identified that coded for both human p38a isoforms (Lee et al. Nature 372, 739). One of the clones which contained the cDNA for p38a-2 (CSBP-2) inserted in the cloning site of pGEX 2T, 3' of the GST coding region was designated pMON 35802. The sequence obtained for this clone is an exact match of the cDNA clone reported by Lee et al. This expression plasmid allows for the production of a GST-p38a fusion protein.

#### Expression of human p38a:

GST/p38a fusion protein was expressed from the plasmid pMON 35802 in *E. coli*, strain DH10B (Life Technologies, Gibco-BRL). Overnight cultures were grown in Luria Broth (LB) containing 100 mg/ml ampicillin. The next day, 500 ml of fresh LB was inoculated with 10 ml of overnight culture, and grown in a 2 liter flask at 37 °C with constant shaking until the culture reached an absorbance of 0.8 at 600 nm. Expression of the fusion protein was induced by addition of isopropyl b-D-thiogalactoside (IPTG) to a final concentration of 0.05 mM. The cultures were shaken for three hours at room temperature, and the cells were harvested by centrifugation. The cell pellets were stored frozen until protein purification.

Purification of p38 Kinase- $\alpha$ :

All chemicals were from Sigma Chemical Co. unless noted. Twenty grams of *E. coli* cell pellet collected from five 1 L shake flask fermentations was resuspended in a volume of PBS (140 mM NaCl, 2.7 mM KCl, 10 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.8 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.3) up to 200 ml. The cell suspension was adjusted to 5 mM DTT with 2 M DTT and then split equally into five 50 ml Falcon conical tubes. The cells were sonicated (Ultrasonics model W375) with a 1 cm probe for 3 X 1 minutes (pulsed) on ice. Lysed cell material was removed by centrifugation (12,000 x g, 15 minutes) and the clarified supernatant applied to glutathione-sepharose resin (Pharmacia).

15 Glutathione-Sephadex Affinity Chromatography:

Twelve ml of a 50% glutathione sepharose-PBS suspension was added to 200 ml clarified supernatant and incubated batchwise for 30 minutes at room temperature. The resin was collected by centrifugation (600 x g, 5 min) and washed with 2 x 150 ml PBS/1% Triton X-100, followed by 4 x 40 ml PBS. To cleave the p38 kinase from the GST-p38 fusion protein, the glutathione-sepharose resin was resuspended in 6 ml PBS containing 250 units thrombin protease (Pharmacia, specific activity > 7500 units/mg) and mixed gently for 4 hours at room temperature. The glutathione-sepharose resin was removed by centrifugation (600 x g, 5 min) and washed 2 x 6 ml with PBS. The PBS wash fractions and digest supernatant containing p38 kinase protein were pooled and adjusted to 0.3 mM PMSF.

Mono Q Anion Exchange Chromatography:

The thrombin-cleaved p38 kinase was further purified by FPLC-anion exchange chromatography. Thrombin-cleaved sample was diluted 2-fold with Buffer A (25 mM HEPES, pH 7.5, 25 mM beta-glycerophosphate, 2 mM DTT, 5% glycerol) and injected onto a Mono Q HR 10/10 (Pharmacia) anion exchange column equilibrated with Buffer A. The column was eluted with a 160 ml 0.1 M-0.6 M NaCl/Buffer A gradient (2 ml/minute flowrate). The p38 kinase peak eluting at 200 mM NaCl was collected and concentrated to 3-4 ml with a Filtron 10 concentrator (Filtron Corp.).

Sephacryl S100 Gel Filtration Chromatography:

The concentrated Mono Q- p38 kinase purified sample was purified by gel filtration chromatography (Pharmacia HiPrep 26/60 Sephacryl S100 column equilibrated with Buffer B (50 mM HEPES, pH 7.5, 50 mM NaCl, 2 mM DTT, 5% glycerol)). Protein was eluted from the column with Buffer B at a 0.5 ml/minute flowrate and protein was detected by absorbance at 280 nm. Fractions containing p38 kinase (detected by SDS-polyacrylamide gel electrophoresis) were pooled and frozen at -80 °C. Typical purified protein yields from 5 L *E. coli* shake flasks fermentations were 35 mg p38 kinase.

**In Vitro Assay**

The ability of compounds to inhibit human p38 kinase alpha was evaluated using two in vitro assay methods. In the first method, activated human p38 kinase alpha phosphorylates a biotinylated substrate, PHAS-I (phosphorylated heat and acid stable protein-insulin inducible), in the presence of gamma <sup>32</sup>P-ATP (<sup>32</sup>P-ATP). PHAS-I was biotinylated prior to the assay and provides a means of capturing the substrate which is phosphorylated during the assay. p38 Kinase was activated by MKK6. Compounds were tested in 10 fold serial dilutions over

the range of 100  $\mu$ M to 0.001  $\mu$ M using 1% DMSO. Each concentration of inhibitor was tested in triplicate.

All reactions were carried out in 96 well polypropylene plates. Each reaction well contained 25 mM HEPES pH 7.5, 10 mM magnesium acetate and 50  $\mu$ M unlabeled ATP. Activation of p38 was required to achieve sufficient signal in the assay. Biotinylated PHAS-I was used at 1-2  $\mu$ g per 50  $\mu$ l reaction volume, with a final concentration of 1.5  $\mu$ M. Activated human p38 kinase alpha was used at 1  $\mu$ g per 50  $\mu$ l reaction volume representing a final concentration of 0.3  $\mu$ M. Gamma  $^{32}$ P-ATP was used to follow the phosphorylation of PHAS-I.  $^{32}$ P-ATP has a specific activity of 3000 Ci/mmol and was used at 1.2  $\mu$ Ci per 50  $\mu$ l reaction volume. The reaction proceeded either for one hour or overnight at 30  $^{\circ}$ C.

Following incubation, 20  $\mu$ l of reaction mixture was transferred to a high capacity streptavidin coated filter plate (SAM-streptavidin-matrix, Promega) prewetted with phosphate buffered saline. The transferred reaction mix was allowed to contact the streptavidin membrane of the Promega plate for 1-2 minutes. Following capture of biotinylated PHAS-I with  $^{32}$ P incorporated, each well was washed to remove unincorporated  $^{32}$ P-ATP three times with 2M NaCl, three washes of 2M NaCl with 1% phosphoric, three washes of distilled water and finally a single wash of 95% ethanol. Filter plates were air dried and 20  $\mu$ l of scintillant was added. The plates were sealed and counted.

A second assay format was also employed that is based on p38 kinase alpha induced phosphorylation of EGFRP (epidermal growth factor receptor peptide, a 21 mer) in the presence of  $^{33}$ P-ATP. Compounds were tested in 10 fold serial dilutions over the range of 100 $\mu$ M to 0.001 $\mu$ M in 1% DMSO. Each concentration of inhibitor was tested in triplicate. Compounds were evaluated in 50 $\mu$ l reaction volumes in the presence of 25 mM Hepes pH 7.5,

10 mM magnesium acetate, 4% glycerol, 0.4% bovine serum albumin, 0.4mM DTT, 50 $\mu$ M unlabeled ATP, 25  $\mu$ g EGFRP (200 $\mu$ M), and 0.05 uCi gamma  $^{33}$ P-ATP. Reactions were initiated by addition of 0.09  $\mu$ g of activated, purified human GST-p38 kinase alpha. Activation was carried out using GST-MKK6 (5:1,p38:MKK6) for one hour at 30 °C in the presence of 50 $\mu$ M ATP. Following incubation for 60 minutes at room temperature, the reaction was stopped by addition of 150 $\mu$ l of AG 1X8 resin in 900 mM sodium formate buffer, pH 3.0 (1 volume resin to 2 volumes buffer). The mixture was mixed three times with pipetting and the resin was allowed to settle. A total of 50 $\mu$ l of clarified solution head volume was transferred from the reaction wells to Microlite-2 plates. 150 $\mu$ l of Microscint 40 was then added to each well of the Microlite plate, and the plate was sealed, mixed, and counted.

#### TNF Cell Assays

20

##### Method of Isolation of Human Peripheral Blood Mononuclear Cells:

Human whole blood was collected in Vacutainer tubes containing EDTA as an anticoagulant. A blood sample (7 ml) was carefully layered over 5 ml PMN Cell Isolation Medium (Robbins Scientific) in a 15 ml round bottom centrifuge tube. The sample was centrifuged at 450-500 x g for 30-35 minutes in a swing out rotor at room temperature. After centrifugation, the top band of cells were removed and washed 3 times with PBS w/o calcium or magnesium. The cells were centrifuged at 400 x g for 10 minutes at room temperature. The cells were resuspended in Macrophage Serum Free Medium (Gibco BRL) at a concentration of 2 million cells/ml.

35

##### LPS Stimulation of Human PBMs:



PBM cells (0.1 ml, 2 million/ml) were co-incubated with 0.1 ml compound (10-0.41  $\mu$ M, final concentration) for 1 hour in flat bottom 96 well microtiter plates. Compounds were dissolved in DMSO initially and diluted in TCM for a final concentration of 0.1% DMSO. LPS (Calbiochem, 20 ng/ml, final concentration) was then added at a volume of 0.010 ml. Cultures were incubated overnight at 37 °C. Supernatants were then removed and tested by ELISA for TNF- $\alpha$  and IL1- $\beta$ . Viability was analyzed using MTS. After 0.1 ml supernatant was collected, 0.020 ml MTS was added to remaining 0.1 ml cells. The cells were incubated at 37 °C for 2-4 hours, then the O.D. was measured at 490-650 nm.

15 Maintenance and Differentiation of the U937 Human Histiocytic Lymphoma Cell Line:

U937 cells (ATCC) were propagated in RPMI 1640 containing 10% fetal bovine serum, 100 IU/ml penicillin, 100  $\mu$ g/ml streptomycin, and 2 mM glutamine (Gibco). Fifty million cells in 100 ml media were induced to terminal monocytic differentiation by 24 hour incubation with 20 ng/ml phorbol 12-myristate 13-acetate (Sigma). The cells were washed by centrifugation (200 x g for 5 min) and resuspended in 100 ml fresh medium. After 24-48 hours, the cells were harvested, centrifuged, and resuspended in culture medium at 2 million cells/ml.

LPS Stimulation of TNF production by U937 Cells:

U937 cells (0.1 ml, 2 million/ml) were incubated with 0.1 ml compound (0.004-50  $\mu$ M, final concentration) for 1 hour in 96 well microtiter plates. Compounds were prepared as 10 mM stock solutions in DMSO and diluted in culture medium to yield a final DMSO concentration of 0.1% in the cell assay. LPS (E coli, 100 ng/ml final concentration) was then added at a volume of 0.02 ml. After 4 hour incubation at 37°C, the amount of TNF- $\alpha$

released in the culture medium was quantitated by ELISA. Inhibitory potency is expressed as IC50 ( $\mu$ M).

#### Rat Assay

5       The efficacy of the novel compounds in blocking the production of TNF also was evaluated using a model based on rats challenged with LPS. Male Harlan Lewis rats [Sprague Dawley Co.] were used in this model. Each rat weighed approximately 300 g and was fasted overnight  
10 prior to testing. Compound administration was typically by oral gavage (although intraperitoneal, subcutaneous and intravenous administration were also used in a few instances) 1 to 24 hours prior to the LPS challenge. Rats were administered 30  $\mu$ g/kg LPS [salmonella typhosa, Sigma Co.] intravenously via the tail vein. Blood was  
15 collected via heart puncture 1 hour after the LPS challenge. Serum samples were stored at -20 °C until quantitative analysis of TNF- $\alpha$  by Enzyme Linked-Immuno-Sorbent Assay ("ELISA") [Biosource]. Additional details  
20 of the assay are set forth in Perretti, M., et al., Br. J. Pharmacol. (1993), 110, 868-874, which is incorporated by reference in this application.

#### Mouse Assay

25

##### Mouse Model Of LPS-Induced TNF Alpha Production:

TNF alpha was induced in 10-12 week old BALB/c female mice by tail vein injection with 100 ng lipopolysaccharide (from S. Typhosa) in 0.2 ml saline.  
30 One hour later mice were bled from the retroorbital sinus and TNF concentrations in serum from clotted blood were quantified by ELISA. Typically, peak levels of serum TNF ranged from 2-6 ng/ml one hour after LPS injection.

The compounds tested were administered to fasted  
35 mice by oral gavage as a suspension in 0.2 ml of 0.5% methylcellulose and 0.025% Tween 20 in water at 1 hour or

6 hours prior to LPS injection. The 1 hour protocol allowed evaluation of compound potency at Cmax plasma levels whereas the 6 hour protocol allowed estimation of compound duration of action. Efficacy was determined at each time point as percent inhibition of serum TNF levels relative to LPS injected mice that received vehicle only.

Induction And Assessment Of Collagen-Induced Arthritis In Mice:

Arthritis was induced in mice according to the procedure set forth in J.M. Stuart, Collagen Autoimmune Arthritis, Annual Rev. Immunol. 2:199 (1984), which is incorporated herein by reference. Specifically, arthritis was induced in 8-12 week old DBA/1 male mice by injection of 50  $\mu$ g of chick type II collagen (CII) (provided by Dr. Marie Griffiths, Univ. of Utah, Salt Lake City, UT) in complete Freund's adjuvant (Sigma) on day 0 at the base of the tail. Injection volume was 100  $\mu$ l. Animals were boosted on day 21 with 50  $\mu$ g of CII in incomplete Freund's adjuvant (100  $\mu$ l volume). Animals were evaluated several times each week for signs of arthritis. Any animal with paw redness or swelling was counted as arthritic. Scoring of arthritic paws was conducted in accordance with the procedure set forth in Wooley et al., Genetic Control of Type II Collagen Induced Arthritis in Mice: Factors Influencing Disease Susceptibility and Evidence for Multiple MHC Associated Gene Control., Trans. Proc., 15:180 (1983). Scoring of severity was carried out using a score of 1-3 for each paw (maximal score of 12/mouse). Animals displaying any redness or swelling of digits or the paw were scored as 1. Gross swelling of the whole paw or deformity was scored as 2. Ankylosis of joints was scored as 3. Animals were evaluated for 8 weeks. 8-10 animals per group were used.

Preparation And Administration Of Compounds:

The compounds tested on mice having collagen-induced arthritis were prepared as a suspension in 0.5% methylcellulose (Sigma, St. Louis, MO), 0.025% Tween 20 (Sigma). The compound suspensions were administered by oral gavage in a volume of 0.1 ml b.i.d. Administration began on day 20 post collagen injection and continued daily until final evaluation on day 56. Scoring of arthritic paws was conducted as set forth above.

Results obtained using the above-described assays are set forth in Table I below. p38 assay and U937 cell assay results are expressed as IC<sub>50</sub> (μM). Mouse-LPS assay results are expressed as percent inhibition.

TABLE I

Example	P38α <sup>1</sup> (μM)	p38α <sup>2</sup> (μM)	U937 (μM)	mLPS (6h @ (30mpk)
1	30.00	13.35	10.00	
2		6.21	10.61	
3		2.55	>10.00	
4		0.23	4.70	54
5	1.98		5.53	
6			10.00	
7		5.48	10.00	
8			10.00	
9	2.44	3.46	0.6474	42
10	7.23	0.4	1.5987	76
11	0.695	10	40	
12	0.941	10	-5	
13	0.86	>10	22	
15	5.9	0.75		32

<sup>1</sup> p38α in vitro results based on PHAS-I assay procedure

<sup>2</sup> p38α in vitro results based on EGFRP assay procedure

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of this invention in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other

active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The active compounds and composition may, for example, be administered orally, intravascularly (IV), intraperitoneally, subcutaneously, intramuscularly (IM) or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, hard or soft capsule, lozenges, dispensable powders, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by injection (IV, IM, subcutaneous or jet) as a composition wherein, for example, saline, dextrose, or water may be used as a suitable carrier. The pH of the composition may be adjusted, if necessary, with suitable acid, base, or buffer. Suitable bulking, dispersing, wetting or suspending agents, including mannitol and PEG 400, may also be included in the composition. A suitable parenteral composition can also include a compound formulated as a sterile solid substance, including lyophilized powder, in injection vials. Aqueous solution can be added to dissolve the compound prior to injection. The amount of therapeutically active compounds that are administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the inflammation or inflammation related disorder, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions

may contain active ingredients in the range of about 0.1 to 1000 mg, preferably in the range of about 7.0 to 350 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between about 0.1 and about 50 mg/kg body weight and most preferably between about 0.5 to 30 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day. In the case of skin conditions, it may be preferable to apply a topical preparation of compounds of this invention to the affected area two to four times a day. For disorders of the eye or other external tissues, e.g., mouth and skin, the formulations are preferably applied as a topical gel, spray, ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in



contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of

5 microcapsules, the encapsulating agent may also function as the membrane. The transdermal patch may include the compound in a suitable solvent system with an adhesive system, such as an acrylic emulsion, and a polyester patch. The oily phase of the emulsions of this invention

10 may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included

15 together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-

20 called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl

25 monostearate, and sodium lauryl sulfate, among others. The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations

30 is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate,

35 propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate,



butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.


Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiinflammatory active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% and particularly about 1.5% w/w. For therapeutic purposes, the active compounds of this combination invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil,

cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

- 5       All patent documents listed herein are incorporated by reference. Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

WHAT IS CLAIMED IS:

1. A compound of Formula I



5 wherein

R¹ is selected from hydrido, alkyl, cycloalkyl, alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene, haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl, mercaptoalkyl, alkylthioalkylene, amino, alkylamino, arylamino, aminoalkyl, alkylaminoalkylene, heterocyclylalkylene, aminocarbonylalkylene, and alkylaminocarbonylalkylene; and

R² is selected from hydrido, alkyl, alkenyl, alkynyl, heterocyclyl, haloalkyl, heterocyclylalkyl, amino, alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy, alkoxycarbonyl, carboxyalkyl, aminocarbonylamino, alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl, alkylsulfonylamino, aminosulfonylamino, alkylaminosulfonylamino, and alkynylamino; wherein the heterocyclyl and heterocyclylalkyl groups are optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl, haloalkyl, amino, cyano, and hydroxy; and

Ar¹ is aryl optionally substituted with one or more radicals independently selected from halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino,

aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl; and

5       HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, 10 cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

2. A compound of Claim 1 wherein

R<sup>1</sup> is selected from hydrido, lower alkyl, lower cycloalkyl, lower cycloalkylalkylene, lower haloalkyl, lower hydroxyalkyl, lower alkenyl, lower alkynyl, lower 5 heterocyclyl, lower aralkyl, lower alkoxyalkyl, lower mercaptoalkyl, lower alkylthioalkylene, amino, lower alkylamino, lower arylamino, lower aminoalkyl, lower alkylaminoalkylene, lower heterocyclylalkylene, lower aminocarbonylalkylene, and lower 10 alkylaminocarbonylalkylene; and

R<sup>2</sup> is selected from hydrido, lower alkyl, lower alkenyl, lower alkynyl, lower haloalkyl, lower heterocyclyl, lower heterocyclylalkylene, amino, lower alkylamino, lower alkynylamino, lower aminoalkyl, lower 15 alkylthio, lower carboxy, lower alkoxycarbonyl, lower carboxyalkyl, lower aminocarbonylamino, lower alkylaminocarbonylamino, lower alkylsulfonyl, lower aminosulfonyl, lower alkylsulfonylamino, lower aminosulfonylamino, and lower alkylaminosulfonylamino, 20 wherein the heterocyclyl and heterocyclylalkyl groups are optionally substituted with one or more radicals independently selected from lower alkylthio, lower

alkylsulfonyl, lower alkylsulfinyl, halo, lower alkyl, lower alkoxy, aryloxy, lower heterocyclyl, lower haloalkyl, amino, and cyano; and

Ar<sup>1</sup> is selected from phenyl, biphenyl, and naphthyl, wherein Ar<sup>1</sup> is optionally substituted with one or more radicals independently selected from lower alkylthio, lower alkylsulfonyl, aminosulfonyl, halo, lower alkyl, lower alkenyl, lower alkynyl, lower alkylsulfinyl, cyano, lower alkoxycarbonyl, aminocarbonyl, formyl, lower alkylcarbonylamino, lower haloalkyl, lower alkoxy, lower alkenyloxy, lower alkylldioxy, amino, lower alkylamino, lower aminoalkyl, arylamino, nitro, and halosulfonyl; and

HetAr<sup>2</sup> is pyridinyl or pyrimidinyl optionally substituted with one or more radicals independently selected from lower alkylthio, lower alkylsulfonyl, lower alkylsulfinyl, halo, lower alkyl, lower heterocyclyl, lower alkoxy, lower aralkoxy, lower haloalkyl, amino, cyano, lower aralkyl, lower alkylamino, lower cycloalkylamino, lower arylamino, lower alkynylamino, and lower aralkylamino; or

a pharmaceutically-acceptable salt or tautomer thereof.

### 3. A Compound of Claim 2 wherein

R<sup>1</sup> is selected from hydrido, methyl, ethyl, isopropyl, tert-butyl, isobutyl, trichloroethyl, pentafluoroethyl, heptafluoropropyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, vinyl, allyl, ethynyl, propargyl, morpholinyl, piperidinyl, piperazinyl, benzyl, phenylethyl, morpholinomethyl, morpholinoethyl, pyrrolidinylmethyl, piperazinylmethyl, piperidinylmethyl, pyridinylmethyl, thienylmethyl, methoxymethyl, ethoxymethyl, amino, methylamino, dimethylamino, phenylamino, methylaminomethyl, dimethylaminomethyl, methylaminoethyl, dimethylaminoethyl, cyclopropyl, cyclopentyl, cyclohexyl,

cyclohexylmethyl, hydroxymethyl, hydroxyethyl, methylthio, and methylthiomethyl; and

$R^2$  is selected from hydrido, methyl, ethyl, propyl, isopropyl, tert-butyl, isobutyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, amino, N-methylamino, N,N-dimethylamino, ethynylamino, propargylamino, piperidinyl, piperazinyl, morpholinomethyl, pyrrolidinylmethyl, piperazinylmethyl, piperidinylmethyl, pyridinylmethyl, thienylmethyl, thiazolylmethyl, oxazolylmethyl, pyrimidinylmethyl, quinolylmethyl, isoquinolinylmethyl, imidazolylmethyl, benzimidazolylmethyl, furylmethyl, pyrazinylmethyl, aminocarbonylamino, methylaminocarbonylamino, dimethylaminocarbonylamino, ethylaminocarbonylamino, diethylaminocarbonylamino, methylsulfonylamino, ethylsulfonylamino, aminosulfonylamino, methylaminosulfonylamino, dimethylaminosulfonylamino, ethylaminosulfonylamino, and diethylaminosulfonylamino; and

$Ar^1$  is selected from phenyl, biphenyl, and naphthyl, wherein  $Ar^1$  is optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, aminosulfonyl, methyl, ethyl, isopropyl, tert-butyl, isobutyl, cyano, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl, methylcarbonylamino, trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl, dichloromethyl, chloromethyl, allyl, vinyl, ethynyl, propargyl, methoxy, ethoxy, propyloxy, n-butoxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, aminomethyl, aminoethyl, N-methyl, N-phenylamino, phenylamino, diphenylamino, nitro, and chlorosulfonyl; and

HetAr<sup>2</sup> is selected from pyridinyl and pyrimidinyl, wherein HetAr<sup>2</sup> is optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, methyl, ethyl, isopropyl, tert-butyl, isobutyl, methoxyl, ethoxyl, phenoxyl, benzoxyl, phenethyl, trifluoromethyl, fluoromethyl, difluoromethyl, amino, benzylamino, propargylamino, cyclopropylamino, cyclobutylamino, cyclopentylamino, and cyano; or  
a pharmaceutically-acceptable salt or tautomer thereof.

4. A compound of Claim 3 wherein  
R<sup>1</sup> is hydrido, methyl, ethyl, hydroxyethyl, propargyl, dimethylaminoethyl or morpholinoethyl; and  
R<sup>2</sup> is selected from hydrido, methyl, ethyl, amino, aminocarbonylamino, methylaminocarbonylamino, methylsulfonylamino, aminosulfonylamino, and methylaminosulfonylamino; and  
Ar<sup>1</sup> is phenyl optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, aminosulfonyl, methyl, ethyl, isopropyl, tert-butyl, isobutyl, cyano, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl, methylcarbonylamino, trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl, dichloromethyl, chloromethyl, methoxy, ethoxy, propyloxy, n-butoxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, aminomethyl, aminoethyl, N-methyl, N-phenylamino, phenylamino, diphenylamino, nitro, and chlorosulfonyl; and

HetAr<sup>2</sup> is optionally substituted with one or more radicals independently selected from methylthio, methylsulfinyl, methylsulfonyl, fluoro, chloro, bromo, methyl, ethyl, isopropyl, tert-butyl, isobutyl, methoxyl, ethoxyl, phenoxyl, benzoxyl, trifluoromethyl,



fluoromethyl, difluoromethyl, amino, propargylamino, and cyano; or

a pharmaceutically-acceptable salt or a tautomer thereof.

5. A compound of Claim 4 wherein

R<sup>1</sup> is hydrido or methyl; and

R<sup>2</sup> is hydrido or methyl; and

Ar<sup>1</sup> is phenyl which is optionally substituted with  
5 one or more radicals independently selected fluoro, chloro, methyl, ethyl, trifluoromethyl, methoxy, ethoxy, dimethylamino, and nitro; and

HetAr<sup>2</sup> is optionally substituted with one or more radicals independently selected from methyl, chloro,  
10 fluoro, and trifluoromethyl; or

a pharmaceutically-acceptable salt or tautomer thereof.

6. A compound of Claim 1 wherein R<sup>2</sup> is hydrido.

7. A compound of Claim 2 wherein R<sup>2</sup> is hydrido.

8. A compound of Claim 3 wherein R<sup>2</sup> is hydrido.

9. A compound of Claim 4 wherein R<sup>2</sup> is hydrido.

10. A compound of Claim 5 wherein R<sup>2</sup> is hydrido.

11. A compound of Claim 1 wherein HetAr<sup>2</sup> is optionally substituted pyridinyl.

12. A compound of Claim 2 wherein HetAr<sup>2</sup> is optionally substituted pyridinyl.

13. A compound of Claim 3 wherein HetAr<sup>2</sup> is optionally substituted pyridinyl.

14. A compound of Claim 4 wherein HetAr<sup>2</sup> is optionally substituted pyridinyl.

15. A compound of Claim 5 wherein HetAr<sup>2</sup> is optionally substituted pyridinyl.

16. A compound of Claim 1 wherein R<sup>2</sup> is hydrido, Ar<sup>1</sup> is optionally substituted phenyl, and HetAr<sup>2</sup> is optionally substituted pyridinyl.

17. A compound of Claim 2 wherein R<sup>2</sup> is hydrido, Ar<sup>1</sup> is optionally substituted phenyl, and HetAr<sup>2</sup> is optionally substituted pyridinyl.

18. A compound of Claim 3 wherein R<sup>2</sup> is hydrido, Ar<sup>1</sup> is optionally substituted phenyl, and HetAr<sup>2</sup> is optionally substituted pyridinyl.

19. A compound of Claim 4 wherein R<sup>2</sup> is hydrido and HetAr<sup>2</sup> is optionally substituted pyridinyl.

20. A compound of Claim 5 wherein R<sup>2</sup> is hydrido and HetAr<sup>2</sup> is optionally substituted pyridinyl.

21. A compound of Claim 4 selected from the compounds, their tautomers and their pharmaceutically acceptable salts, of the group consisting of

4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine;

5 4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine;

N-[4(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide;


N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylsulfamide;

10 [4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]urea;

[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]sulfamide;

- 4-(4-chlorophenyl)-1-methyl-3-(4-pyridinyl)-1H-pyrazol-5-amine;
- N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylurea;
- 5 4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine;
- 4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine;
- 4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazole-1-ethanol;
- 4-(4-fluorophenyl)-N,N-dimethyl-3-(4-pyridinyl)-1H-pyrazole-1-ethanamine;
- 10 4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine;
- 4-[4-(4-chlorophenyl)-1H-pyrazol-3-yl]pyridine;
- 1-methyl-4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine; and
- 15 1-methyl-4-[2-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine.

22. A compound of Formula I



wherein

- R¹ is selected from hydrido, lower alkyl, lower cycloalkyl, lower cycloalkylalkylene, lower haloalkyl, lower hydroxyalkyl, lower alkynyl, lower aralkyl, lower alkoxyalkyl, lower mercaptoalkyl, lower alkylthioalkylene, amino, lower alkylamino, lower arylamino, lower aminoalkyl, lower alkylaminoalkylene, lower heterocyclalkylene, lower aminocarbonylalkylene, and lower alkylaminocarbonylalkylene; and
- 5
- 10

$R^2$  is selected from hydrido, lower alkyl, lower cycloalkyl, lower cycloalkenyl, aryl selected from phenyl and biphenyl, lower aralkyl, lower haloalkyl, lower heterocyclyl, lower heterocyclylalkylene, amino, lower arylamino, lower alkylamino, lower alkynylamino, lower aminocarbonylamino, lower alkylaminocarbonylamino, sulfonyl, lower alkylsulfonylamino, lower aminosulfonylamino, and lower alkylaminosulfonylamino, wherein the cycloalkyl, aryl and heterocyclyl groups are optionally substituted with one or more radicals independently selected from lower alkylthio, lower alkylsulfonyl, lower alkylsulfinyl, halo, lower alkyl, lower alkoxy, aryloxy, lower heterocyclyl, lower haloalkyl, amino, and cyano; and

$Ar^1$  is phenyl optionally substituted with one or more radicals independently selected from halo, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, amino, aminocarbonyl, cyano, lower alkoxycarbonyl, formyl, aminosulfonyl, lower alkylamino, nitro, lower arylamino, lower alkylcarbonylamino, halosulfonyl, lower aminoalkyl, and lower haloalkyl; and

$HetAr^2$  is pyridinyl optionally substituted with one or more radicals independently selected from lower alkylthio, lower alkylsulfonyl, lower alkylsulfinyl, halo, lower alkyl, lower heterocyclyl, lower alkoxy, lower aralkoxy, lower haloalkyl, amino, cyano, lower aralkyl, lower alkylamino, lower cycloalkylamino, lower aralkylamino, and lower arylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

23. A compound of claim 22 wherein

$R^1$  is hydrido, methyl, hydroxyethyl, dimethylaminoethyl, propargyl, or morpholinoethyl; and

R<sup>2</sup> is selected from hydrido, methyl, ethyl, amino, aminocarbonylamino, methylaminocarbonylamino, methylsulfonylamino, aminosulfonylamino, and methylaminosulfonylamino;

24. A compound of claim 22 wherein R<sup>2</sup> is hydrido.

25. A compound of claim 22 wherein Ar<sup>1</sup> is phenyl substituted with one or more halogen radicals.

26. A compound of claim 22 wherein R<sup>2</sup> is hydrido and Ar<sup>1</sup> is phenyl substituted with one or more halogen radicals.

27. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from the compounds of Claim 1; or a pharmaceutically salt or tautomer thereof.

28. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from the compounds of Claim 2; or a pharmaceutically salt or tautomer thereof.

29. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from the compounds of Claim 3; or a pharmaceutically salt or tautomer thereof.


30. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from the compounds of Claim 4; or a pharmaceutically salt or tautomer thereof.

31. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said

compound selected from the compounds of Claim 5; or a pharmaceutically salt or tautomer thereof.

32. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from the compounds of Claim 21; or a pharmaceutically salt or tautomer thereof.

33. A method of treating a TNF mediated disorder, said method comprising treating the subject having or susceptible to such disorder with a therapeutically-effective amount of a compound of Formula I



wherein

$R^1$  is selected from hydrido, alkyl, cycloalkyl, alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene, haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl, mercaptoalkyl, alkylthioalkylene, amino, alkylamino, arylamino, aminoalkyl, alkylaminoalkylene, heterocyclylalkylene, aminocarbonylalkylene, and alkylaminocarbonylalkylene; and

$R^2$  is selected from hydrido, alkyl, alkenyl, alkynyl, heterocyclyl, haloalkyl, heterocyclylalkyl, amino, alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy, alkoxycarbonyl, carboxyalkyl, aminocarbonylamino, alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl, alkylsulfonylamino, aminosulfonylamino, alkylaminosulfonylamino, and alkynylamino; wherein the

heterocyclyl and heterocyclylalkyl groups are optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl, haloalkyl, amino, cyano, and hydroxy; and


Ar<sup>1</sup> is aryl optionally substituted with one or more radicals independently selected from halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino, aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl; and

HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

34. A method of treating a p38 kinase mediated disorder, said method comprising treating the subject having or susceptible to such disorder with a therapeutically-effective amount of a compound of Formula

I






wherein

- R<sup>1</sup> is selected from hydrido, alkyl, cycloalkyl, alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene, haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl, mercaptoalkyl, alkylthioalkylene, amino, alkylamino, arylamino, aminoalkyl, alkylaminoalkylene, heterocyclylalkylene, aminocarbonylalkylene, and alkylaminocarbonylalkylene; and
- R<sup>2</sup> is selected from hydrido, alkyl, alkenyl, alkynyl, heterocyclyl, haloalkyl, heterocyclylalkyl, amino, alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy, alkoxycarbonyl, carboxyalkyl, aminocarbonylamino, alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl, alkylsulfonylamino, aminosulfonylamino, alkylaminosulfonylamino, and alkynylamino; wherein the heterocyclyl and heterocyclylalkyl groups are optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl, haloalkyl, amino, cyano, and hydroxy; and
- Ar<sup>1</sup> is aryl optionally substituted with one or more radicals independently selected from halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkyldioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino, aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl; and
- HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

35. A method of treating inflammation, said method comprising treating the subject having or susceptible to such condition with a therapeutically-effective amount of a compound of Formula I



wherein


- $R^1$  is selected from hydrido, alkyl, cycloalkyl, alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene, haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl, mercaptoalkyl, alkylthioalkylene, amino, alkylamino, arylamino, aminoalkyl, alkylaminoalkylene, heterocyclylalkylene, aminocarbonylalkylene, and alkylaminocarbonylalkylene; and
- $R^2$  is selected from hydrido, alkyl, alkenyl, alkynyl, heterocyclyl, haloalkyl, heterocyclylalkyl, amino, alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy, alkoxy carbonyl, carboxyalkyl, aminocarbonylamino, alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl, alkylsulfonylamino, aminosulfonylamino, alkylaminosulfonylamino, and alkynylamino; wherein the heterocyclyl and heterocyclylalkyl groups are optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl, haloalkyl, amino, cyano, and hydroxy; and

Ar<sup>1</sup> is aryl optionally substituted with one or more radicals independently selected from halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino, aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl; and

HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer thereof.

36. A method of treating arthritis, said method comprising treating the subject having or susceptible to such condition with a therapeutically-effective amount of a compound of Formula I



wherein

R<sup>1</sup> is selected from hydrido, alkyl, cycloalkyl, alkenyl, alkynyl, heterocyclyl, cycloalkylalkylene, haloalkyl, hydroxyalkyl, aralkyl, alkoxyalkyl, mercaptoalkyl, alkylthioalkylene, amino, alkylamino,

arylamino, aminoalkyl, alkylaminoalkylene,  
heterocyclylalkylene, aminocarbonylalkylene, and  
alkylaminocarbonylalkylene; and

$R^2$  is selected from hydrido, alkyl, alkenyl, alkynyl,  
5 heterocyclyl, haloalkyl, heterocyclylalkyl, amino,  
alkylamino, aminoalkyl, alkoxy, alkylthio, carboxy,  
alkoxycarbonyl, carboxyalkyl, aminocarbonylamino,  
alkylaminocarbonylamino, alkylsulfonyl, aminosulfonyl,  
alkylsulfonylamino, aminosulfonylamino,  
10 alkylaminosulfonylamino, and alkynylamino; wherein the  
heterocyclyl and heterocyclylalkyl groups are optionally  
substituted with one or more radicals independently  
selected from alkylthio, alkylsulfonyl, alkylsulfinyl,  
halo, alkyl, alkoxy, aryloxy, aralkoxy, heterocyclyl,  
15 haloalkyl, amino, cyano, and hydroxy; and

$Ar^1$  is aryl optionally substituted with one or more  
radicals independently selected from halo, alkyl,  
alkenyl, alkynyl, alkoxy, alkenoxy, alkylldioxy,  
alkylthio, alkylsulfinyl, alkylsulfonyl, amino,  
20 aminocarbonyl, cyano, alkoxycarbonyl, formyl,  
aminosulfonyl, alkylamino, nitro, arylamino,  
alkylcarbonylamino, halosulfonyl, aminoalkyl, and  
haloalkyl; and

$HetAr^2$  is pyridinyl, pyrimidinyl or quinolinyl  
25 optionally substituted with one or more radicals  
independently selected from alkylthio, alkylsulfonyl,  
alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy,  
aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino,  
cycloalkylamino, cycloalkenylamino, arylamino,  
30 alkynylamino, and aralkylamino; or

a pharmaceutically-acceptable salt or a tautomer  
thereof.

37. The method of Claim 33 wherein the TNF mediated  
disorder is selected from the group of disorders  
consisting of bone resorption, graft vs. host reaction,

atherosclerosis, arthritis, osteoarthritis, rheumatoid arthritis, gout, psoriasis, topical inflammatory disease state, adult respiratory distress syndrome, asthma, chronic pulmonary inflammatory disease, cardiac  
5 reperfusion injury, renal reperfusion injury, thrombus, glomerulonephritis, Crohn's disease, ulcerative colitis, inflammatory bowel disease and cachexia.

38. The method of Claim 33 wherein the TNF mediated disorder is inflammation.

39. The method of Claim 33 wherein the TNF mediated disorder is arthritis.

40. The method of Claim 33 wherein the TNF mediated disorder is asthma.

41. The method of claim 33 wherein the compound is selected from the compounds, their tautomers and their pharmaceutically acceptable salts, of the group consisting of  
5 4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine;  
4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine;  
N-[4(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide;  
N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-  
10 N'-methylsulfamide;  
[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]urea;  
[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]sulfamide;  
4-(4-chlorophenyl)-1-methyl-3-(4-pyridinyl)-1H-pyrazol-5-  
15 amine;  
N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylurea;  
4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine;  
4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine;

4- (4-fluorophenyl) -3- (4-pyridinyl) -1H-pyrazole-1-ethanol;  
4- (4-fluorophenyl) -N,N-dimethyl-3- (4-pyridinyl) -1H-  
pyrazole-1-ethanamine;  
4- [2- [4- (4-fluorophenyl) -3- (4-pyridinyl) -1H-pyrazol-1-  
5 yl]ethyl]morpholine;  
4- [4- (4-chlorophenyl) -1H-pyrazol-3-yl]pyridine;  
1-methyl-4- [2- [4- (4-fluorophenyl) -3- (4-pyridinyl) -1H-  
pyrazol-1-yl]]piperidine; and  
1-methyl-4- [2- [4- (4-fluorophenyl) -5- (4-pyridinyl) -1H-  
10 pyrazol-1-yl]]piperidine.

42. The method of Claim 34 wherein the disorder is a p38 $\alpha$  kinase mediated disorder.

43. The method of Claim 34 wherein the P38 kinase mediated disorder is selected from the group of disorders consisting of bone resorption, graft vs. host reaction, atherosclerosis, arthritis, osteoarthritis, rheumatoid  
5 arthritis, gout, psoriasis, topical inflammatory disease state, adult respiratory distress syndrome, asthma, chronic pulmonary inflammatory disease, cardiac reperfusion injury, renal reperfusion injury, thrombus, glomerulonephritis, Crohn's disease, ulcerative colitis,  
10 inflammatory bowel disease and cachexia.

44. The method of Claim 34 wherein the p38 kinase mediated disorder is inflammation.

45. The method of Claim 34 wherein the p38 kinase mediated disorder is arthritis.

46. The method of Claim 34 wherein the p38 kinase mediated disorder is asthma.

47. The method of claim 34 wherein the compound is selected from the compounds, their tautomers and their

pharmaceutically acceptable salts, of the group consisting of

4-(3-methyl-4-phenyl-1H-pyrazol-5-yl)pyridine;  
4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-amine;  
5 N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]methanesulfonamide;  
N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylsulfamide;  
[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]urea;  
10 [4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]sulfamide;  
4-(4-chlorophenyl)-1-methyl-3-(4-pyridinyl)-1H-pyrazol-5-amine;  
N-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-3-yl]-N'-methylurea;  
15 4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine;  
4-[4-(4-fluorophenyl)-1-methyl-1H-pyrazol-3-yl]pyridine;  
4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazole-1-ethanol;  
4-(4-fluorophenyl)-N,N-dimethyl-3-(4-pyridinyl)-1H-pyrazole-1-ethanamine;  
20 4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]ethyl]morpholine;  
4-[4-(4-chlorophenyl)-1H-pyrazol-3-yl]pyridine;  
1-methyl-4-[2-[4-(4-fluorophenyl)-3-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine; and  
25 1-methyl-4-[2-[4-(4-fluorophenyl)-5-(4-pyridinyl)-1H-pyrazol-1-yl]]piperidine.

48. A compound of claim 1 that is 4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine or a pharmaceutically-acceptable salt or a tautomer thereof.


49. The method of claim 33 wherein the compound is 4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine or a pharmaceutically-acceptable salt or a tautomer thereof.



50. The method of claim 34 wherein the compound is 4-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]pyridine or a pharmaceutically-acceptable salt or a tautomer thereof.



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07D 401/04, 401/14, A61K 31/415, 31/44</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 98/52937</b> <b>(43) International Publication Date:</b> 26 November 1998 (26.11.98)
<b>(21) International Application Number:</b> PCT/US98/10807 <b>(22) International Filing Date:</b> 22 May 1998 (22.05.98)  <b>(30) Priority Data:</b> 60/047,535                      22 May 1997 (22.05.97)                      US  <b>(71) Applicant (for all designated States except US):</b> G.D. SEARLE AND CO. [US/US]; P.O. Box 5110, Chicago, IL 60680 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ANANTANARAYAN, Ashok [US/US]; 54 Lisk Drive, Hainesville, IL 60015 (US). CLARE, Michael [GB/US]; 5154 West Brown Street, Skokie, IL 60077 (US). GENG, Lifeng [CN/US]; 5300 Davis Street, Skokie, IL 60077 (US). HANSON, Gunnar, J. [US/US]; 7410 Keystone Avenue, Skokie, IL 60076 (US). PARTIS, Richard, A. [US/US]; 2221 Noyes Street, Evanston, IL 60012 (US). STEALEY, Michael, A. [US/US]; 502 Juniper Parkway, Libertyville, IL 60048 (US). WEIER, Richard, M. [US/US]; 240 Hickory Court, Lake Bluff, IL 60044 (US).		<b>(74) Agents:</b> WILLIAMS, Scott, A. et al.; Senniger, Powers, Leavitt & Roedel, 16th floor F1, One Metropolitan Square, St. Louis, MO 63102 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 11 March 1999 (11.03.99)
<b>(54) Title:</b> 4-ARYL-3(5)-HETEROARYL SUBSTITUTED PYRAZOLES AS P38 KINASE  <b>(57) Abstract</b> <p>A class of pyrazole derivatives is described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula (I), wherein R<sup>1</sup> and R<sup>2</sup> are as described in the specification; and Ar<sup>1</sup> is aryl optionally substituted with one or more radicals independently selected from halo, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkylidioxy, alkylthio, alkylsulfinyl, alkylsulfonyl, amino, aminocarbonyl, cyano, alkoxycarbonyl, formyl, aminosulfonyl, alkylamino, nitro, arylamino, alkylcarbonylamino, halosulfonyl, aminoalkyl, and haloalkyl; and HetAr<sup>2</sup> is pyridinyl, pyrimidinyl or quinolinyl optionally substituted with one or more radicals independently selected from alkylthio, alkylsulfonyl, alkylsulfinyl, halo, alkyl, heterocyclyl, alkoxy, aralkoxy, haloalkyl, amino, cyano, aralkyl, alkylamino, cycloalkylamino, cycloalkenylamino, arylamino, alkynylamino, and aralkylamino; or a pharmaceutically-acceptable salt or a tautomer thereof.</p> <div style="text-align: right;">  <p style="text-align: right;">(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/US 98/10807

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D401/04 C07D401/14 A61K31/415 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>JP 05 017 470 A (OTSUKA PHARMACEUT FACTORY INC) 26 January 1993 cited in the application</p> <p>see pages 9-17; the examples (or compound no.) 1-19, 21-23, 31, 35 and 37 &amp; DATABASE WPI Section Ch, Week 9309 Derwent Publications Ltd., London, GB; Class B03, AN 93-071109 see abstract</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>1-4, 11-14, 21-23, 27-30, 32, 35, 36</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 September 1998

Date of mailing of the international search report

25.09.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Fink, D

## INTERNATIONAL SEARCH REPORT

Inter. .onal Application No

PCT/US 98/10807

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI  Section Ch, Week 9638  Derwent Publications Ltd., London, GB;  Class B02, AN 96-379287  XP002076468  -&amp; JP 08 183 787 A (EISAI CO LTD)  see abstract  &amp; JP 08 183 787 A (EISAI CO. LTD.) 16 July  1996</p> <p>---</p>	1-32,48
X	<p>EP 0 418 845 A (FUJISAWA PHARMACEUTICAL  CO) 27 March 1991  cited in the application</p> <p>see page 55; claim 1  see page 21, line 54 - page 22, line 12</p> <p>---</p>	1-3,6-8, 11-13, 16-18, 21, 27-29, 35,36,48
X	<p>EP 0 515 041 A (DU PONT) 25 November 1992  cited in the application  see page 196 - page 197; claim 1  see page 200 - page 201; claim 8</p> <p>---</p>	1-26,48
X	<p>MOERCK R E ET AL: "Pyrazole addition to  2H-azirines. Structure of purported 2:1  adducts of 2H-azirines with  sym-tetrazines"  J. CHEM. SOC., CHEM. COMMUN. ,  no. 19, 1974,  pages 782-783, XP002076466  see page 783, column 1, the scheme, the  compound (2c)</p> <p>---</p>	1,2,11, 12,22
X	<p>DANNHARDT G ET AL: "Transformations of  pyrrolidine enaminones yielding  (.omega.-aminopropyl)pyrazoles"  ARCH. PHARM. ,  vol. 321, no. 1, 1988,  pages 17-19, XP002076467  see page 17, column 1, the compounds 3f  and 3g</p> <p>---</p>	1,2,11, 12,27,28
X	<p>WO 97 01551 A (FUJISAWA PHARMACEUTICAL CO  ;AKAHANE ATSUSHI (JP); KURODA SATORU (J)  16 January 1997  cited in the application  see page 72; claim 1</p> <p>---</p>	1,11,27

-/--

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/10807

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 559 137 A (ADAMS JERRY L ET AL) 24 September 1996 cited in the application	1,2,6,7, 11,12, 16,17, 27,28, 33,35-40
Y	see column 2, line 66 - column 5, line 29 see column 2, line 66 - column 5, line 29 ---	1-50
Y	WO 96 21452 A (ADAMS JERRY LEROY ;BOEHM JEFFREY CHARLES (US); GALLAGHER TIMOTHY F) 18 July 1996 see the whole document; in particular page 61, example 23; and page 41, lines 22-24 ---	1-50
A	EP 0 531 901 A (FUJISAWA PHARMACEUTICAL CO) 17 March 1993 see the whole document -----	1-50

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 98/10807

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 33-47, 49 and 50 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Appl. Application No

PCT/US 98/10807

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 05017470 A	26-01-1993	JP 2753659 B	20-05-1998
EP 0418845 A	27-03-1991	AT 126216 T	15-08-1995
		AU 637142 B	20-05-1993
		AU 6307290 A	18-04-1991
		CA 2025599 A	23-03-1991
		CN 1050382 A	03-04-1991
		DE 69021472 D	14-09-1995
		DE 69021472 T	25-01-1996
		DK 418845 T	18-09-1995
		ES 2088933 T	01-10-1996
		GR 3017100 T	30-11-1995
		HU 9500344 A	28-09-1995
		IE 68857 B	24-07-1996
		IL 95675 A	31-03-1996
		JP 2586713 B	05-03-1997
		JP 3141261 A	17-06-1991
		NO 301006 B	01-09-1997
		PT 95389 A	22-05-1991
		RU 2021990 C	30-10-1994
		RU 2059622 C	10-05-1996
		US 5134142 A	28-07-1992
EP 0515041 A	25-11-1992	AU 1988692 A	21-12-1992
		CN 1066069 A	11-11-1992
		WO 9219615 A	12-11-1992
WO 9701551 A	16-01-1997	NONE	
US 5559137 A	24-09-1996	JP 10500413 T	13-01-1998
		WO 9531451 A	23-11-1995
WO 9621452 A	18-07-1996	US 5593992 A	14-01-1997
		AU 4657296 A	31-07-1996
		BG 101727 A	31-03-1998
		BR 9606904 A	21-10-1997
		CA 2209938 A	18-07-1996
		CN 1177299 A	25-03-1998
		CZ 9702158 A	15-04-1998
		EP 0809499 A	03-12-1997

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. onal Application No

PCT/US 98/10807

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9621452 A		FI 972901 A	08-09-1997
		NO 973167 A	08-09-1997
		PL 322249 A	19-01-1998
		SK 90297 A	14-01-1998
		US 5670527 A	23-09-1997
		US 5593991 A	14-01-1997
		US 5663334 A	02-09-1997
		ZA 9600094 A	24-07-1996
<hr/>			
EP 0531901 A	17-03-1993	AU 2280592 A	11-03-1993
		CA 2077732 A	10-03-1993
		CN 1070404 A	31-03-1993
		HU 65204 A	02-05-1994
		JP 7252256 A	03-10-1995
		JP 2057962 C	10-06-1996
		JP 6287188 A	11-10-1994
		JP 7088386 B	27-09-1995
		MX 9205145 A	01-04-1993
		US 5478827 A	26-12-1995
		US 5624931 A	29-04-1997
		US 5356897 A	18-10-1994
		ZA 9206417 A	15-03-1993
		CN 1075965 A	08-09-1993
<hr/>			